

A DIXMIER-DOUADY THEORY FOR STRONGLY SELF-ABSORBING C^* -ALGEBRAS

MARIUS DADARLAT AND ULRICH PENNIG

ABSTRACT. We show that the automorphism group $\text{Aut}(A)$ of a strongly self-absorbing C^* -algebra A is contractible and that the automorphism group $\text{Aut}(A \otimes \mathbb{K})$ of the stabilization of A has the homotopy type of a CW-complex whose homotopy groups we compute. We extend the classification of locally trivial bundles of C^* -algebras with compact operators \mathbb{K} as fibers by Dixmier and Douady to the case where the fibers are isomorphic to a stabilized strongly self-absorbing C^* -algebra $A \otimes \mathbb{K}$.

We prove that the classifying space for these bundles $B\text{Aut}(A \otimes \mathbb{K})$ has the structure of an infinite loop space with respect to the tensor product. Thus, bundles with fibers $A \otimes \mathbb{K}$ are classified by a generalized continuous cohomology theory which is computable via the Atiyah-Hirzebruch spectral sequence. This allows us to introduce rational characteristic classes for such bundles. A necessary and sufficient K -theoretical condition for local triviality is given for continuous fields with fibers $A \otimes \mathbb{K}$ over spaces of finite covering dimension.

CONTENTS

1. Introduction	1
2. The topology of $\text{Aut}(A \otimes \mathbb{K})$ for strongly self-absorbing algebras	4
2.1. Strongly self-absorbing C^* -algebras	4
2.2. Contractibility of $\text{Aut}(A)$	5
2.3. The homotopy type of $\text{Aut}_0(A \otimes \mathbb{K})$	7
2.4. The homotopy type of $\text{Aut}(A \otimes \mathbb{K})$	9
2.5. The topological group $\text{Aut}(A \otimes \mathbb{K})$ is well-pointed	13
3. The infinite loop space structure of $B\text{Aut}(A \otimes \mathbb{K})$	14
3.1. Permutative categories and infinite loop spaces	14
3.2. The tensor product of $A \otimes \mathbb{K}$ -bundles	16
4. A generalized Dixmier-Douady theory	20
References	24

1. INTRODUCTION

Continuous field C^* -algebras are employed as versatile tools in several areas, including index and representation theory, the Novikov and the Baum-Connes conjectures [11], (deformation) quantization [37, 27] and the study of the quantum Hall effect [5]. While continuous fields play the

M.D. was partially supported by NSF grant #DMS-1101305.

role of bundles in C^* -algebra theory, the underlying bundle structure is typically not locally trivial. Nevertheless, these bundles have sufficient continuity properties to allow for local propagation of interesting K -theory invariants along their fibers. Continuous field C^* -algebras with simple fibers occur naturally as the class of C^* -algebras with Hausdorff primitive spectrum.

In 1963 Dixmier and Douady published their classic study of continuous field C^* -algebras with fibers stably isomorphic to the compact operators $\mathbb{K} = \mathbb{K}(H)$ (H an infinite dimensional Hilbert space) over a paracompact base space X [17]. Dixmier and Douady proved that:

Theorem 1.1. *The isomorphism classes of locally trivial fields over X with fibers \mathbb{K} form a group under the operation of tensor product and this group is isomorphic to $\check{H}^3(X, \mathbb{Z})$.*

Theorem 1.2. *If X is finite dimensional, then a separable continuous field B over X with fibers isomorphic to \mathbb{K} is locally trivial if and only if it satisfies Fell's condition, i.e. each point of X has a closed neighborhood V such that the restriction of B to V contains a projection of constant rank 1.*

The corresponding characteristic class $\delta(B) \in \check{H}^3(X, \mathbb{Z})$ is now known as the Dixmier-Douady invariant. Most prominent among its applications is its appearance as twisting class in twisted K -theory [18, 40, 3, 4], due to which it also gained importance in string theory as the D -brane charge [45, 8]. A recent friendly introduction to the Dixmier-Douady theory can be found in [41].

The present paper extends the results of Dixmier and Douady to continuous field C^* -algebras which have stabilized strongly self-absorbing C^* -algebras $A \otimes \mathbb{K}$ as fibers. The class of strongly self-absorbing C^* -algebras, introduced by Toms and Winter [49], is closed under tensor products and contains C^* -algebras that are cornerstones of Elliott's classification program of simple nuclear C^* -algebras: the Cuntz algebras \mathcal{O}_2 and \mathcal{O}_∞ , the Jiang-Su algebra \mathcal{Z} , the canonical anticommutation relations algebra $M_{2\infty}$ and in fact all UHF-algebras of infinite type. These are separable C^* -algebras singled out by a crucial property: there exists an isomorphism $A \rightarrow A \otimes A$, which is unitarily homotopic to the map $a \mapsto a \otimes 1_A$, [16, 50]. Using this property, which is equivalent to, but formally much stronger than the original definition of [49], we prove that

- $\text{Aut}(A)$ is contractible in the point-norm topology.
- $\text{Aut}(A \otimes \mathbb{K})$ is well-pointed and it has the homotopy type of a CW-complex.
- The classifying space $B\text{Aut}(A \otimes \mathbb{K})$ of locally trivial C^* -algebra bundles with fiber $A \otimes \mathbb{K}$ carries an H -space structure induced by the tensor product. Moreover, this tensor product multiplication is homotopy commutative up to all higher homotopies and therefore equips $B\text{Aut}(A \otimes \mathbb{K})$ with the structure of an infinite loop space by results of Segal and May.

These properties mirror entirely the corresponding properties of $\text{Aut}(\mathbb{K}) = PU(H)$ and $B\text{Aut}(\mathbb{K}) = BPU(H)$ obtained by their identification with the Eilenberg-MacLane spaces $K(\mathbb{Z}, 2)$ and respectively $K(\mathbb{Z}, 3)$ which we implicitly reprove as they correspond to the case $A = \mathbb{C}$. Recall that if X is paracompact Hausdorff, then $\check{H}^n(X, \mathbb{Z}) \cong [X, K(\mathbb{Z}, n)]$, [24].

It is worth noting that while the obstructions to having a natural group structure on the isomorphism classes of locally trivial continuous fields with fiber $A \otimes \mathbb{K}$ – such as nontrivial Samelson products [36, sec.6] – do vanish in the strongly self-absorbing case, that is not necessarily true for

general self-absorbing C^* -algebras, i.e. those with $A \otimes A \cong A$. This motivates yet again our choice of fibers. In complete analogy with Theorem 1.1 we have:

Theorem A. *Let X be a compact metrizable space and let A be a strongly self-absorbing C^* -algebra. The set $\text{Bun}_X(A \otimes \mathbb{K})$ of isomorphism classes of locally trivial fields over X with fiber $A \otimes \mathbb{K}$ becomes an abelian group under the operation of tensor product. Moreover, this group is isomorphic to $E_A^1(X)$, the first group of a generalized connective cohomology theory $E_A^*(X)$ defined by the infinite loop space $B\text{Aut}(A \otimes \mathbb{K})$.*

We also show that the zero group $E_A^0(X)$ computes the homotopy classes $[X, \text{Aut}(A \otimes \mathbb{K})]$ and it is isomorphic to the group of positive invertible elements of the abelian ring $K_0(C(X) \otimes A)$, denoted by $K_0(C(X) \otimes A)_+^\times$, for $A \neq \mathbb{C}$. In particular, we fully compute the coefficients of $E_A^*(X)$, as they are given by the homotopy groups

$$\pi_i(\text{Aut}(A \otimes \mathbb{K})) = \begin{cases} K_0(A)_+^\times & \text{if } i = 0 \\ K_i(A) & \text{if } i \geq 1. \end{cases}$$

$K_0(A)$ has a natural ring structure with unit given by the class of 1_A . $K_0(A)^\times$ denotes the group of multiplicative elements of $K_0(A)$ and $K_0(A)_+^\times$ is its subgroup consisting of positive elements.

The Atiyah-Hirzebruch spectral sequence then allows us to obtain classification results for locally trivial $A \otimes \mathbb{K}$ -bundles over X . In the case of the universal UHF algebra $M_{\mathbb{Q}}$, bundles with fiber $M_{\mathbb{Q}} \otimes \mathbb{K}$ are essentially classified by the ordinary rational cohomology groups of odd degree of the underlying space:

$$\text{Bun}_X(M_{\mathbb{Q}} \otimes \mathbb{K}) \cong E_{M_{\mathbb{Q}}}^1(X) \cong H^1(X, \mathbb{Q}_+^\times) \oplus \bigoplus_{k \geq 1} H^{2k+1}(X, \mathbb{Q}).$$

A similar result holds for bundles with fiber $\mathcal{O}_\infty \otimes M_{\mathbb{Q}} \otimes \mathbb{K}$, see Corollary 4.4. It follows that if A is any strongly self-absorbing C^* -algebra that satisfies the UCT, then there are rational characteristic classes $\delta_k : \text{Bun}_X(A \otimes \mathbb{K}) \rightarrow H^{2k+1}(X, \mathbb{Q})$ such that $\delta_k(B_1 \otimes B_2) = \delta_k(B_1) + \delta_k(B_2)$.

An unexpected consequence of our results is that for any strongly self-absorbing C^* -algebra A , if two bundles $B_1, B_2 \in \text{Bun}_X(A \otimes \mathbb{K})$ become isomorphic after tensoring with \mathcal{O}_∞ , then they must be isomorphic in the first place, see Corollary 4.8.

Our result concerning local triviality is the following generalization of Theorem 1.2 which involves a K -theoretic interpretation of Fell's condition.

Theorem B. *Let X be a locally compact metrizable space of finite covering dimension and let A be a strongly self-absorbing C^* -algebra. A separable continuous field B over X with fibers abstractly isomorphic to $A \otimes \mathbb{K}$ is locally trivial if and only if for each point $x \in X$, there exist a closed neighborhood V of x and a projection $p \in B(V)$ such that $[p(v)] \in K_0(B(v))^\times$ for all $v \in V$.*

The condition that X is finite dimensional is essential in Theorem B, as shown by examples constructed in [17] for $A = \mathbb{C}$, [23] for $A = M_{\mathbb{Q}}$ and [14] for $A = \mathcal{O}_2$.

Let us recall that a C^* -algebra isomorphic to the compact operators on a separable (possibly finite dimensional) Hilbert space is called an elementary C^* -algebra. Dixmier and Douady gave two other results concerning continuous fields of elementary C^* -algebras:

(i) If B is a continuous field of elementary C^* -algebras that satisfies Fell's condition, then $B \otimes \mathbb{K}$ is locally trivial.

(ii) The class $\delta(B) \in \check{H}^3(X, \mathbb{Z})$ can be defined for any continuous field of elementary C^* -algebras that satisfies Fell's condition. Moreover B is isomorphic to the compact operators of a continuous field of Hilbert spaces if and only if $\delta(B) = 0$.

We extend (i) and the first part of (ii) to general strongly self-absorbing C^* -algebras, but we must require finite dimensionality for either the fiber or the base space in order to obtain a perfect analogy with these results, see Corollaries 4.9 and 4.10. These restrictions are necessary. Indeed, while any unital separable continuous field C^* -algebra with fiber \mathbb{C} over X is locally trivial (in fact isomorphic to $C_0(X)$), automatic local triviality fails if \mathbb{C} is replaced by strongly self-absorbing C^* -algebras such as $M_{\mathbb{Q}}$ and \mathcal{O}_2 , see [23] and [14].

This fact also explains why the second part of (ii) is specific to fields of elementary C^* -algebras. Our set-up allows us to associate rational characteristic classes to any continuous fields (satisfying Fell's condition) whose fibers are Morita equivalent to strongly self-absorbing C^* -algebras which are not necessarily mutually isomorphic. Such fields are typically very far from being locally trivial. We refer the reader to Section 4 for further discussion.

The homotopy equivalence $\text{Aut}(A \otimes \mathbb{K}) \simeq K_0(A)_+^{\times} \times BU(A)$ (see Corollary 2.17) suggests that the generalized cohomology theory associated to $\text{Aut}(A \otimes \mathbb{K})$ is very closely related to the unit spectrum $GL_1(KU^A)$ of topological K -theory with coefficients in the group $K_0(A)$. This is again parallel to the Dixmier-Douady theory, where we have $\text{Aut}(\mathbb{K}) = PU(H) \simeq BU(1) \subset GL_1(KU)$. We will make this connection precise elsewhere. Let us just mention here that the homotopy equivalence $\text{Aut}(\mathcal{Z} \otimes \mathbb{K}) \simeq BU$ deloops to a homotopy equivalence $B\text{Aut}(\mathcal{Z} \otimes \mathbb{K}) \simeq B(BU_{\otimes})$. This unveils a very natural operator algebra realization of the classic Ω -spectrum $B(BU_{\otimes})$ introduced by Segal [44].

The authors are grateful to Johannes Ebert, Peter May and Jim McClure for a number of useful discussions.

2. THE TOPOLOGY OF $\text{Aut}(A \otimes \mathbb{K})$ FOR STRONGLY SELF-ABSORBING ALGEBRAS

The automorphism group $\text{Aut}(B)$ of a separable C^* -algebra B , equipped with the point-norm topology, is a separable and metrizable topological group. In particular its topology is compactly generated. We are going to show in this section that if A is a strongly self-absorbing C^* -algebra, then $\text{Aut}(A \otimes \mathbb{K})$ is well-pointed and has the homotopy type of a CW -complex. This will enable us to apply the standard techniques of algebraic topology, in particular when it comes to dealing with its classifying space. We denote by \simeq the relation of homotopy equivalence.

2.1. Strongly self-absorbing C^* -algebras. Let us recall from [49] that a C^* -algebra A is strongly self-absorbing if it is separable, unital and there exists a $*$ -isomorphism $\psi : A \rightarrow A \otimes A$ such that ψ is approximately unitarily equivalent to the map $l : A \rightarrow A \otimes A$, $l(a) = a \otimes 1_A$. It

follows from [16] and [50] that ψ and l must be in fact unitarily homotopy equivalent, see Theorem 2.1(b). Note that, unlike [49], we don't exclude the complex numbers \mathbb{C} from the class of strongly self-absorbing C^* -algebras. For future reference, we collect under one roof an important series of results due to several authors.

Theorem 2.1. *A strongly self-absorbing C^* -algebra A has the following properties:*

- (a) A is simple, nuclear and is either stably finite or purely infinite; if it is stably finite, then it admits a unique trace, see [49] and references therein.
- (b) Let B be a unital separable C^* -algebra. For any two unital $*$ -homomorphisms $\alpha, \beta : A \rightarrow B \otimes A$ there is a continuous path of unitaries $(u_t)_{t \in [0,1]}$ in $B \otimes A$ such that $u_0 = 1$ and $\lim_{t \rightarrow 1} \|u_t \alpha(a) u_t^* - \beta(a)\| = 0$ for all $a \in A$. This property was proved in [16, Thm.2.2] under the assumption that A is K_1 -injective. Winter [50] has shown that any infinite dimensional strongly self-absorbing C^* -algebra A is \mathcal{Z} -stable, i.e. $A \otimes \mathcal{Z} \cong A$, and hence A is K_1 -injective by a result of Rørdam [39].
- (c) Any unital \mathcal{Z} -stable C^* -algebra has cancellation of full projections by a result of Jiang [25, Thm.1]. In particular, if B is a separable unital C^* -algebra and $A \neq \mathbb{C}$, then $B \otimes A$ is isomorphic to $B \otimes A \otimes \mathcal{Z}$ and hence it has cancellation of full projections.
- (d) If B is a unital \mathcal{Z} -stable C^* -algebra, then $\pi_0(U(B)) \cong K_1(B)$, by [25, Thm.2].
- (e) If A satisfies the Universal Coefficient Theorem (UCT) in KK-theory, then $K_1(A) = 0$ by [49]. If in addition A is purely infinite, then A is isomorphic to either \mathcal{O}_2 or \mathcal{O}_∞ or a tensor product of \mathcal{O}_∞ with a UHF-algebra of infinite type [49, Cor.5.2].

Notation. For C^* -algebras A, B we denote by $\text{Hom}(A, B)$ the space of full $*$ -homomorphisms from A to B and by $\text{End}(A)$ the space of full $*$ -endomorphisms of A . Recall that a $*$ -homomorphism $\varphi : A \rightarrow B$ is full if for any nonzero element $a \in A$, the closed ideal generated by $\varphi(a)$ is equal to B . If A is a unital C^* -algebra, we denote by $K_0(A)_+$ the subsemigroup of $K_0(A)$ consisting of classes $[p]$ of full projections $p \in A \otimes \mathbb{K}$.

2.2. Contractibility of $\text{Aut}(A)$. While it is known from [16, Cor.3.1] that $\text{Aut}(A)$ is weakly contractible in the point norm-topology, we can strengthen this result by combining it with the idea of half-flips from [49].

Let B be a separable C^* -algebra and let $e \in B$ be a projection. Consider the following spaces of $*$ -endomorphisms of B endowed with the point-norm topology.

$$\text{End}_e(B) = \{\alpha \in \text{End}(B) : \alpha(e) = e\}, \quad \text{Aut}_e(B) = \{\alpha \in \text{Aut}(B) : \alpha(e) = e\}.$$

Let $l, r : B \rightarrow B \otimes B$ (minimal tensor product) be defined by $l(b) = b \otimes e$ and $r(b) = e \otimes b$.

Lemma 2.2. *Suppose that there is a continuous map $\psi : [0, 1] \rightarrow \text{Hom}(B, B \otimes B)$ such that $\psi(0) = l$, $\psi(1) = r$, $\psi(t)(e) = e \otimes e$ and $\psi(t)$ is a $*$ -isomorphism for all $t \in (0, 1)$. Then $\text{Aut}_e(B)$ and $\text{End}_e(B)$ are contractible spaces.*

Proof. First we deal with $\text{Aut}_e(B)$. Consider $H: I \times \text{Aut}_e(B) \rightarrow \text{Aut}_e(B)$ defined by

$$(1) \quad H(t, \alpha) = \begin{cases} \alpha & \text{for } t = 0 \\ \psi(t)^{-1} \circ (\alpha \otimes \text{id}_B) \circ \psi(t) & \text{for } 0 < t < 1, \\ \text{id}_B & \text{for } t = 1. \end{cases}$$

Note that $H(t, \alpha)(e) = e$ since $\psi(t)(e) = e \otimes e$. Observe that $(\alpha \otimes \text{id}_B) \circ l = l \circ \alpha$. It is straightforward to verify the continuity of H at points (α, t) with $t \neq 0$ and $t \neq 1$. Let $b \in B$, let $t_n \in (0, 1)$ be a net converging to 0 and let $\alpha_i \in \text{Aut}_e(B)$ be a net converging to $\alpha \in \text{Aut}_e(B)$. The estimate,

$$\begin{aligned} & \|(\psi(t_n)^{-1} \circ (\alpha_i \otimes \text{id}_B) \circ \psi(t_n))(b) - \alpha(b)\| = \|(\alpha_i \otimes \text{id}_B) \circ \psi(t_n)(b) - \psi(t_n) \circ \alpha(b)\| \\ & \leq \|(\alpha_i \otimes \text{id}_B) \circ \psi(t_n)(b) - (\alpha_i \otimes \text{id}_B) \circ l(b)\| + \|(\alpha_i \otimes \text{id}_B) \circ l(b) - (\alpha \otimes \text{id}_B) \circ l(b)\| \\ & + \|(\alpha \otimes \text{id}_B) \circ l(b) - \psi(t_n) \circ \alpha(b)\| \\ & \leq \|\psi(t_n)(b) - l(b)\| + \|\alpha_i(b) - \alpha(b)\| + \|(\alpha \otimes \text{id}_B) \circ l(b) - \psi(t_n) \circ \alpha(b)\| \end{aligned}$$

implies the continuity of H at $(\alpha, 0)$. An analogous argument using $(\alpha \otimes \text{id}_B) \circ r = r$ shows continuity at $(\alpha, 1)$. We also have $H(t, \text{id}_B) = \text{id}_B$ for all $t \in [0, 1]$. Thus, H provides a (strong) deformation retraction of $\text{Aut}_e(B)$ to id_B . The argument for the contractibility of $\text{End}_e(B)$ is entirely similar. One observes that equation (1) also defines a map $H: I \times \text{End}_e(B) \rightarrow \text{End}_e(B)$ which gives a deformation retraction of $\text{End}_e(B)$ to id_B . \square

Theorem 2.3. *Let A be a strongly self-absorbing C^* -algebra. Then $\text{Aut}(A)$ and $\text{End}_{1_A}(A)$ are contractible spaces.*

Proof. Let $l, r: A \rightarrow A \otimes A$ be the maps $l(a) = a \otimes 1_A$ and $r(a) = 1_A \otimes a$. Fix an isomorphism $\psi: A \rightarrow A \otimes A$. It follows from Theorem 2.1(b) that there exists a continuous path of unitaries $u: (0, 1] \rightarrow U(A \otimes A)$ with $u(1) = 1_{A \otimes A}$ such that

$$\lim_{t \rightarrow 0} \|u(t) \psi(a) u(t)^* - l(a)\| = 0.$$

Define $\psi_l: (0, 1] \rightarrow \text{Iso}(A, A \otimes A)$ by $\psi_l(t) = \text{Ad}_{u(t)} \circ \psi$. Likewise there is a continuous path of unitaries $v: [0, 1) \rightarrow U(A \otimes A)$ with $v(0) = 1_{A \otimes A}$ and such that

$$\lim_{t \rightarrow 1} \|v(t) \psi(a) v(t)^* - r(a)\| = 0.$$

Define $\psi_r: [0, 1) \rightarrow \text{Iso}(A, A \otimes A)$ by $\psi_r(t) = \text{Ad}_{v(t)} \circ \psi$. By juxtaposing the paths ψ_l and ψ_r we obtain a homotopy from l to r which satisfies the assumptions of Lemma 2.2 with $e = 1_A$. It follows that $\text{Aut}(A)$ and $\text{End}_{1_A}(A)$ are contractible spaces. \square

The following is a minor variation of a result of Blackadar and Hermann and Rosenberg.

Lemma 2.4. *Let A and B be separable AF -algebras and let $e \in A$ be a projection. Suppose that $\varphi, \psi: A \rightarrow B$ are two $*$ -homomorphisms such that $\varphi(e) = \psi(e)$ and $\varphi_* = \psi_*: K_0(A) \rightarrow K_0(B)$. Then there is a continuous map $u: [0, 1) \rightarrow U(B^+)$ with $u(0) = 1$, $[u(t), \psi(e)] = 0$ for all $t \in [0, 1)$ and such that $\lim_{t \rightarrow 1} \|u(t) \psi(a) u(t)^* - \varphi(a)\| = 0$ for all $a \in A$.*

Proof. If B is a nonunital C^* -algebra, we regard B as a C^* -subalgebra of its unitization B^+ . Write A as the closure of an increasing union of finite dimensional C^* -subalgebras $A_n \subset A_{n+1}$ with $A_0 = \mathbb{C}e$. Since $\phi_* = \psi_*$, for each $n \geq 0$ we find a unitary $u_n \in U(B^+)$ such that $u_n \psi(x) u_n^* = \varphi(x)$ for all $x \in A_n$ and $u_0 = 1$. Observe that $w_n = u_{n+1}^* u_n$ is a unitary in the commutant C_n of $\psi(A_n)$ in B^+ . This commutant is known to be an AF-algebra, see [21, Lemma 3.1]. Therefore there is a continuous path of unitaries $t \mapsto W_n(t) \in U(C_n)$, $t \in [n, n+1]$, such that $W_n(n) = w_n$ and $W_n(n+1) = 1$. Define a continuous map $u : [0, \infty) \rightarrow U(B)$ by $u(t) = u_{n+1} W_n(t)$, $t \in [n, n+1]$. One verifies immediately that $[u(t), \psi(e)] = 0$ for all t and that $u(t) \psi(x) u(t)^* = \varphi(x)$ for all $x \in A_n$ and $t \in [n, n+1]$. It follows that $\lim_{t \rightarrow \infty} \|u(t) \psi(a) u(t)^* - \varphi(a)\| = 0$ for all $a \in A$. \square

Theorem 2.5. *Let A be a strongly self-absorbing C^* -algebra and let $e \in \mathbb{K}$ be a rank-1 projection. Then the stabilizer group $\text{Aut}_{1 \otimes e}(A \otimes \mathbb{K})$ and the space $\text{End}_{1 \otimes e}(A \otimes \mathbb{K})$ are contractible.*

Proof. We shall use the following consequence of Lemma 2.4. Let $\varphi_0, \varphi_1 : \mathbb{K} \rightarrow \mathbb{K} \otimes \mathbb{K}$ be two $*$ -homomorphisms such that $\varphi_0(e) = \varphi_1(e) = e \otimes e$. Fix a $*$ -isomorphism $\psi_{1/2} : \mathbb{K} \rightarrow \mathbb{K} \otimes \mathbb{K}$ with $\psi_{1/2}(e) = e \otimes e$. By applying Lemma 2.4 to both pairs $(\varphi_i, \psi_{1/2})$, $i = 0, 1$, we find a continuous map $\psi : [0, 1] \rightarrow \text{Hom}(\mathbb{K}, \mathbb{K} \otimes \mathbb{K})$ such that $\psi(0) = \varphi_0$, $\psi(1) = \varphi_1$, $\psi(t)(e) = e \otimes e$ and $\psi(t)$ is a $*$ -isomorphism for all $t \in (0, 1)$.

We proceed in much the same way as the proof of Theorem 2.3, by applying Lemma 2.2. Let $l, r : A \rightarrow A \otimes A$ be defined by $l(a) = a \otimes 1_A$ and $r(a) = 1_A \otimes a$. We have seen in the proof of Theorem 2.3 that there is a continuous map $\psi : [0, 1] \rightarrow \text{Hom}(A, A \otimes A)$ such that $\psi(0) = l$, $\psi(1) = r$, and $\psi(t)$ is a $*$ -isomorphism for all $t \in (0, 1)$.

Let $l_{\mathbb{K}}, r_{\mathbb{K}} : \mathbb{K} \rightarrow \mathbb{K} \otimes \mathbb{K}$ be given by $l_{\mathbb{K}}(x) = x \otimes e$, $r_{\mathbb{K}}(x) = e \otimes x$. Using the remark from the beginning of the proof, we find a continuous map $\psi_{\mathbb{K}} : [0, 1] \rightarrow \text{Hom}(\mathbb{K}, \mathbb{K} \otimes \mathbb{K})$ such that $\psi_{\mathbb{K}}(0) = l_{\mathbb{K}}$, $\psi_{\mathbb{K}}(1) = r_{\mathbb{K}}$, $\psi_{\mathbb{K}}(t)(e) = e \otimes e$ and $\psi_{\mathbb{K}}(t)$ is a $*$ -isomorphism for all $t \in (0, 1)$.

Let $A_{\mathbb{K}} = A \otimes \mathbb{K}$ and consider the $*$ -homomorphisms $\hat{l}, \hat{r} : A_{\mathbb{K}} \rightarrow A_{\mathbb{K}} \otimes A_{\mathbb{K}}$ with

$$\hat{l} = \sigma \circ (l \otimes l_{\mathbb{K}}) \quad \text{and} \quad \hat{r} = \sigma \circ (r \otimes r_{\mathbb{K}}),$$

where $\sigma : A \otimes (A \otimes \mathbb{K}) \otimes \mathbb{K} \rightarrow (A \otimes \mathbb{K}) \otimes (A \otimes \mathbb{K})$ interchanges the second and third tensor factor. Note that $\hat{l}(a \otimes x) = (a \otimes x) \otimes (1_A \otimes e)$ and $\hat{r}(a \otimes x) = (1_A \otimes e) \otimes (a \otimes x)$ for $a \in A$ and $x \in \mathbb{K}$. Define $\hat{\psi} : [0, 1] \rightarrow \text{Hom}(A_{\mathbb{K}}, A_{\mathbb{K}} \otimes A_{\mathbb{K}})$ by $\hat{\psi} = \sigma \circ (\psi \otimes \psi_{\mathbb{K}})$. Then $\hat{\psi}(0) = \hat{l}$, $\hat{\psi}(1) = \hat{r}$, $\hat{\psi}(t)(1_A \otimes e) = (1_A \otimes e) \otimes (1_A \otimes e)$ and $\hat{\psi}(t)$ is an isomorphism for all $t \in (0, 1)$. It follows by Lemma 2.2 that $\text{Aut}_{1 \otimes e}(A \otimes \mathbb{K})$ and $\text{End}_{1 \otimes e}(A \otimes \mathbb{K})$ are contractible. \square

Remark 2.6. Taking $A = \mathbb{C}$, Thm. 2.5 reproves the contractibility of $U(H)$ in the strong topology.

2.3. The homotopy type of $\text{Aut}_0(A \otimes \mathbb{K})$. For a C^* -algebra B we denote by $\text{Aut}_0(B)$ the path-connected component of the identity. We have seen in Theorem 2.3 that for a strongly self-absorbing C^* -algebra A the space $\text{Aut}(A)$ is contractible. In particular, it has the homotopy type of a CW-complex. In this and the following subsections, we will extend the latter statement to $\text{Aut}(A \otimes \mathbb{K})$, which is no longer contractible, but has a very interesting homotopy type. We start by considering the subspace of projections in $A \otimes \mathbb{K}$, denoted by $\text{Pr}(A \otimes \mathbb{K})$.

Lemma 2.7. *Let B be a C^* -algebra. The space $\text{Pr}(B)$ has the homotopy type of a CW-complex.*

Proof. Let B_{sa} be the real Banach space of self-adjoint elements in B . Consider the subset U of B_{sa} consisting of all elements which do not have $1/2$ in the spectrum. Since invertibility is an open condition, U is an open subset of B_{sa} and therefore has the homotopy type of a CW-complex by [28, Cor.5.5, p.134]. Since $\sigma(p) \subset \{0, 1\}$ for any projection $p \in B$, we have $\mathcal{P}r(B) \subset U$. Let f be the characteristic function of the interval $(\frac{1}{2}, \infty)$. By functional calculus, f induces a continuous map $U \rightarrow \mathcal{P}r(B)$, $a \mapsto f(a)$, which restricts to the identity on $\mathcal{P}r(B)$. Thus, $\mathcal{P}r(B)$ is dominated by a space having the homotopy type of a CW-complex. By [28, Cor.3.9, p.127] it is homotopy equivalent to a CW-complex itself. \square

Let e be a rank-1 projection in \mathbb{K} . We define $\mathcal{P}r_0(A \otimes \mathbb{K})$ to be the connected component of $1 \otimes e \in \mathcal{P}r(A \otimes \mathbb{K})$. It does not depend on the choice of e as long as the rank of e is equal to 1.

Lemma 2.8. *Let A be a unital C^* -algebra and let $e \in \mathbb{K}$ be a rank-1 projection. Then the sequence $\text{Aut}_{1 \otimes e}(A \otimes \mathbb{K}) \rightarrow \text{Aut}_0(A \otimes \mathbb{K}) \rightarrow \mathcal{P}r_0(A \otimes \mathbb{K})$ where the first map is the inclusion and the second sends α to $\alpha(1 \otimes e)$ is a locally trivial fiber bundle over a paracompact base space and therefore a Hurewicz fibration.*

Proof. This is a particular case of a more general result. Let B be a C^* -algebra, let $q \in \mathcal{P}r(B)$ and let $\mathcal{P}r_0(B)$ be the path-component of q . Then $\pi : \text{Aut}_0(B) \rightarrow \mathcal{P}r_0(B)$, $\pi(\alpha) = \alpha(q)$ is in fact a principal fiber bundle with structure group $\text{Aut}_q(B)$. The map π is well-defined. Indeed, if α is homotopic to id_B , then the projection $\alpha(q)$ is connected to q by a continuous path in $\mathcal{P}r(B)$.

Let $U_0(B^+)$ denote the path-component of 1 in the unitary group of the unitization of B . Thus, for $u \in U_0(B^+)$ we have $\text{Ad}_u \in \text{Aut}_0(B)$. By definition any $p \in \mathcal{P}r_0(B)$ is homotopic to q . Therefore p and q are also unitarily equivalent via a unitary $u \in U_0(B^+)$. Since $\pi(\text{Ad}_u) = p$ it follows that π is surjective. Let $p_0 \in \mathcal{P}r_0(B)$ and let U be its open neighborhood given by $U = \{p \in \mathcal{P}r_0(B) \mid \|p - p_0\| < 1\}$. If $p \in U$, then $x_p = p_0 p + (1 - p_0)(1 - p)$ is an invertible element of B^+ . It follows that $u_p = x_p(x_p^* x_p)^{-\frac{1}{2}}$ is a unitary in $U_0(B^+)$ and the map $p \mapsto u_p$ is continuous with respect to the norm topologies [7, Prop.II.3.3.4]. Choose a unitary $v \in U_0(B^+)$ such that $p_0 = v q v^*$. Then $\sigma_{p_0} : U \rightarrow \text{Aut}_0(B)$, $p \mapsto \text{Ad}_{u_p^* v}$ is a continuous section of π over U . This completes the proof. \square

Corollary 2.9. *Let A be a strongly self-absorbing C^* -algebra. Then $\text{Aut}_0(A \otimes \mathbb{K})$ has the homotopy type of a CW-complex, which is homotopy equivalent to $\mathcal{P}r_0(A \otimes \mathbb{K}) \simeq BU(A)$.*

Proof. By Lemma 2.8 and Theorem 2.5 the space $\text{Aut}_0(A \otimes \mathbb{K})$ is the total space of a fibration where both base and fiber have the homotopy type of a CW-complex. Now the statement follows from [42, Thm.2], Theorem 2.5, except that it remains to argue that $\mathcal{P}r_0(A \otimes \mathbb{K}) \simeq BU(A)$. This is certainly known. The group $U(M(A \otimes \mathbb{K}))$ acts continuously and transitively on $\mathcal{P}r_0(A \otimes \mathbb{K})$ via $u \mapsto u(1 \otimes e)u^*$ with stabilizer $U(A) \times U(M(A \otimes \mathbb{K}))$. By the contractibility of $U(M(A \otimes \mathbb{K}))$ [12],

$$U(A) \rightarrow U(M(A \otimes \mathbb{K}))/1_A \times U(M(A \otimes \mathbb{K})) \rightarrow \mathcal{P}r_0(A \otimes \mathbb{K})$$

is the *universal* principal $U(A)$ -bundle. One uses the map $p \mapsto u_p$ constructed in the proof of Lemma 2.8 in order to verify local triviality. Thus $\mathcal{P}r_0(A \otimes \mathbb{K})$ is a model for $BU(A)$. \square

2.4. The homotopy type of $\text{Aut}(A \otimes \mathbb{K})$. In this section we compute the homotopy classes $[X, \text{End}(A \otimes \mathbb{K})]$ and $[X, \text{Aut}(A \otimes \mathbb{K})]$ in the case of a strongly self-absorbing C^* -algebra A and a compact metrizable space X , see Theorem 2.22. A similar topic was studied for Kirchberg algebras in [13]. Throughout this subsection e is a rank-1 projection in \mathbb{K} . Given a unital ring R , we denote by R^\times the group of units in R . It is easily seen that $K_0(C(X) \otimes A)$ carries a ring structure with multiplication induced by an isomorphism $\psi : A \otimes \mathbb{K} \rightarrow A \otimes \mathbb{K} \otimes A \otimes \mathbb{K}$ which maps $1_A \otimes e$ to $1_A \otimes e \otimes 1_A \otimes e$. This structure does not depend on the choice ψ by Theorem 2.5. We identify the space of continuous maps from X to $\text{End}(A \otimes \mathbb{K})$ with $\text{Hom}(A \otimes \mathbb{K}, C(X) \otimes A \otimes \mathbb{K})$ and with $\text{End}_{C(X)}(C(X) \otimes A \otimes \mathbb{K})$. Similarly, we will identify the space of continuous maps from X to $\text{Aut}(A \otimes \mathbb{K})$ with $\text{Aut}_{C(X)}(C(X) \otimes A \otimes \mathbb{K})$.

Lemma 2.10. *Let A and B be unital separable C^* -algebras. Suppose that $p \in B \otimes \mathbb{K}$ is a full projection such that there is a unital $*$ -homomorphism $\theta : A \rightarrow p(B \otimes \mathbb{K})p$. Then there is a $*$ -homomorphism $\varphi : A \otimes \mathbb{K} \rightarrow B \otimes \mathbb{K}$ such that $\varphi(1 \otimes e) = p$. If θ is an isomorphism, then we can choose φ to be an isomorphism.*

Proof. We denote by \sim Murray-von Neumann equivalence of projections. Let us recall that if $q, r \in B \otimes \mathbb{K}$ are projections, then $q \sim r$ if and only if there is $u \in U(M(B \otimes \mathbb{K}))$ such that $uqu^* = r$, [34, Lemma 1.10]. Since p is a full projection in $B \otimes \mathbb{K}$, by [9], there is $v \in M(B \otimes \mathbb{K} \otimes \mathbb{K})$ such that $v^*v = p \otimes I$ and $vv^* = 1 \otimes I \otimes I$. Then $\gamma : p(B \otimes \mathbb{K})p \otimes \mathbb{K} \rightarrow B \otimes \mathbb{K} \otimes \mathbb{K}$, $\gamma(a) = vav^*$, is an isomorphism with the property that $\gamma(p \otimes e) = v(p \otimes e)v^* \sim (p \otimes e)(v^*v)(p \otimes e) = p \otimes e$. The map $\mathbb{K} \rightarrow \mathbb{K} \otimes \mathbb{K}$, $x \mapsto x \otimes e$ is homotopic to a $*$ -isomorphism as observed in the proof of Theorem 2.5. It follows that the map $B \otimes \mathbb{K} \rightarrow B \otimes \mathbb{K} \otimes \mathbb{K}$, $b \otimes x \mapsto b \otimes x \otimes e$ is also homotopic to a $*$ -isomorphism μ . Note that $\mu(p) \sim p \otimes e \sim \gamma(p \otimes e)$. Thus, after conjugating μ by a unitary in $M(B \otimes \mathbb{K})$ we may arrange that $\mu(p) = \gamma(p \otimes e)$. It follows that $\varphi = \mu^{-1} \circ \gamma \circ (\theta \otimes \text{id}_{\mathbb{K}}) \in \text{Hom}(A \otimes \mathbb{K}, B \otimes \mathbb{K})$ has the property that $\varphi(1 \otimes e) = p$. Finally note that if θ is an isomorphism then so is φ . \square

Corollary 2.11 (Kodaka, [26]). *Let A be a separable unital C^* -algebra and let $p \in A \otimes \mathbb{K}$ be a full projection. Then $p(A \otimes \mathbb{K})p \cong A$ if and only if there is $\alpha \in \text{Aut}(A \otimes \mathbb{K})$ such that $\alpha(1 \otimes e) = p$.*

Proposition 2.12. *Let A be a strongly self-absorbing C^* -algebra and let B be a separable unital C^* -algebra such that $B \cong B \otimes A$. Let $\varphi, \psi : A \otimes \mathbb{K} \rightarrow B \otimes \mathbb{K}$ be two full $*$ -homomorphisms. Suppose that $[\varphi(1_A \otimes e)] = [\psi(1_A \otimes e)] \in K_0(B)$. Then (i) φ is homotopic to ψ and (ii) φ is approximately unitarily equivalent to ψ , written $\varphi \approx_u \psi$.*

Proof. (i) For C^* -algebras A, B we denote by $[A, B]_\#$ the homotopy classes of full $*$ -homomorphisms $\varphi : A \rightarrow B$. The inclusion $A \cong A \otimes e \hookrightarrow A \otimes \mathbb{K}$ induces a restriction map $\rho : [A \otimes \mathbb{K}, B \otimes \mathbb{K}]_\# \rightarrow [A, B \otimes \mathbb{K}]_\#$. Thomsen showed that ρ is bijective, see [48, Lemma 1.4]. Since the map $[\varphi] \mapsto [\varphi(1 \otimes e)]$ factors through ρ , it suffices to show that the map $[A, B \otimes \mathbb{K}]_\# \rightarrow K_0(B)$, $\varphi \mapsto [\varphi(1)]$ is injective. Let $\varphi, \psi : A \rightarrow B \otimes \mathbb{K}$ be two full $*$ -homomorphisms. Suppose that $[\varphi(1)] = [\psi(1)]$. Since B has cancellation of full projections by Theorem 2.1(c), after conjugation by a unitary in the contractible group $U(M(B \otimes \mathbb{K}))$, we may assume that $\varphi(1) = \psi(1) = p \in \text{Pr}(B \otimes \mathbb{K})$. The C^* -algebra $p(B \otimes \mathbb{K})p$ is A -absorbing by [49, Cor.3.1]. It follows that the $*$ -homomorphisms $\varphi, \psi : A \rightarrow p(B \otimes \mathbb{K})p$ are homotopic by Theorem 2.1(b).

(ii) It suffices to prove approximate unitary equivalence for the restrictions of φ and ψ to $A \otimes M_n(\mathbb{C})$ for any $n \geq 1$. Let (e_{ij}) denote the canonical matrix unit of $M_n(\mathbb{C})$, $p_{ij} = \varphi(1 \otimes e_{ij})$, $q_{ij} = \psi(1 \otimes e_{ij})$, $p_n = \varphi(1 \otimes 1_n)$ and $q_n = \psi(1 \otimes 1_n)$. By reasoning as in part (a), we find a partial isometry $v \in B \otimes \mathbb{K}$ such that $v^*v = p_{11}$ and $vv^* = q_{11}$. By [34, Lemma 1.10] there is a partial isometry $w \in M(B \otimes \mathbb{K})$ such that $w^*w = 1 - p_n$ and $ww^* = 1 - q_n$. It follows that $V = w + \sum_{k=1}^n q_{k1}vp_{1k}$ is a unitary in $M(A \otimes \mathbb{K})$ such that $V\varphi(1 \otimes x)V^* = \psi(1 \otimes x)$ for all $x \in M_n(\mathbb{C})$. Thus after conjugating φ by a unitary we may assume that $\varphi(1 \otimes x) = \psi(1 \otimes x)$ for all $x \in M_n(\mathbb{C})$. Let us observe that if $a \in A$ and $u \in U(p_{11}(B \otimes \mathbb{K})p_{11})$, then $U = (1 - p_n) + \sum_{k=1}^n p_{k1}up_{1k} \in U(M(B \otimes \mathbb{K}))$ satisfies $U\varphi(a \otimes e_{ij})U^* - \psi(a \otimes e_{ij}) = p_{i1}(u\varphi(a \otimes e_{11})u^* - \psi(a \otimes e_{11}))p_{1j}$. This reduces our task to proving approximate unitary equivalence for the unital maps $A \otimes e \rightarrow p(B \otimes \mathbb{K})p$ induced by φ and ψ , where $p = \varphi(1 \otimes e)$. Since $p(B \otimes \mathbb{K})p$ is A -absorbing, this follows from Theorem 2.1(b). \square

Next we consider the case when $B = C(X) \otimes A$ in Lemma 2.10. Next we compare two natural multiplicative H -space structures on $\text{End}(A \otimes \mathbb{K})$.

Lemma 2.13. *Let X be a topological space, let A be a strongly self-absorbing C^* -algebra and let $\psi: A \otimes \mathbb{K} \rightarrow (A \otimes \mathbb{K}) \otimes (A \otimes \mathbb{K})$ be a $*$ -isomorphism. The two operations $*$ and \circ on $G = [X, \text{End}(A \otimes \mathbb{K})]$ defined by*

$$[\alpha] * [\beta] = [\psi^{-1} \circ (\alpha \otimes \beta) \circ \psi] \quad \text{and} \quad [\alpha] \circ [\beta] = [\alpha \circ \beta],$$

where $\alpha \otimes \beta: X \rightarrow \text{End}((A \otimes \mathbb{K})^{\otimes 2})$ denotes the pointwise tensor product, agree and are both associative and commutative. Moreover, $*$ does not depend on the choice of ψ and is a group operation when restricted to $\text{Aut}(A \otimes \mathbb{K})$.

Proof. First, let $\hat{\psi}$, \hat{l} and \hat{r} be as in the proof of Theorem 2.5 and use $\psi = \hat{\psi}(\frac{1}{2})$ in the definition of the operation $*$. Given α, β, δ and $\gamma \in C(X, \text{End}(A \otimes \mathbb{K}))$ we have

$$\begin{aligned} ([\alpha] * [\beta]) \circ ([\gamma] * [\delta]) &= [\psi^{-1} \circ (\alpha \otimes \beta) \circ \psi \circ \psi^{-1} \circ (\gamma \otimes \delta) \circ \psi] \\ &= [\psi^{-1} \circ ((\alpha \circ \gamma) \otimes (\beta \circ \delta)) \circ \psi] = ([\alpha] \circ [\gamma]) * ([\beta] \circ [\delta]). \end{aligned}$$

Thus, the Eckmann-Hilton [19] argument will imply that $*$ and \circ agree and are both associative and commutative for this particular choice of ψ if we can show that $\text{id}_{A \otimes \mathbb{K}}$ is a unit for the operation $*$. Just as in the proof of Theorem 2.5 we can see that $\hat{\psi}(t/2)^{-1} \circ (\alpha \otimes \text{id}_{A \otimes \mathbb{K}}) \circ \hat{\psi}(t/2)$, $t \in [0, 1]$, is a homotopy from α to $\psi^{-1} \circ (\alpha \otimes \text{id}_{A \otimes \mathbb{K}}) \circ \psi$ with respect to the point-norm topology on $\text{End}(A \otimes \mathbb{K})$ proving that $\text{id}_{A \otimes \mathbb{K}}$ is a right unit. The analogous argument for $\hat{\psi}((t+1)/2)^{-1} \circ (\text{id}_{A \otimes \mathbb{K}} \otimes \alpha) \circ \hat{\psi}((t+1)/2)$ shows that $\text{id}_{A \otimes \mathbb{K}}$ is also a left unit.

If ψ is chosen arbitrarily, we have $\psi = \hat{\psi}(\frac{1}{2}) \circ \kappa$ for some $\kappa \in \text{Aut}(A \otimes \mathbb{K})$. We denote the corresponding operations by $*_{\psi}$ and $*_{\hat{\psi}}$ and have

$$[\alpha] *_{\psi} [\beta] = [\kappa^{-1} \circ \hat{\psi}^{-1}(\frac{1}{2}) \circ (\alpha \otimes \beta) \circ \hat{\psi}(\frac{1}{2}) \circ \kappa] = [\kappa^{-1}] \circ ([\alpha] *_{\hat{\psi}} [\beta]) \circ [\kappa] = [\alpha] *_{\hat{\psi}} [\beta]$$

by the homotopy commutativity of \circ . This proves the independence of $*$ from the choice of the isomorphism ψ . \square

We denote by \approx_u the relation of approximate unitary equivalence for $*$ -homomorphisms.

Lemma 2.14. *Let A be a strongly self-absorbing C^* -algebra. If $p \in A \otimes \mathbb{K}$ is a nonzero projection, the following conditions are equivalent:*

- (i) $p(A \otimes \mathbb{K})p \cong A$
- (ii) *There is $\alpha \in \text{Aut}(A \otimes \mathbb{K})$ such that $\alpha(1 \otimes e) = p$.*
- (iii) $[p] \in K_0(A)_+^\times$

We denote by $\text{Pr}(A \otimes \mathbb{K})^\times$ the set of all projections satisfying these equivalent conditions.

Proof. (i) \Leftrightarrow (ii) This follows from Corollary 2.11.

(ii) \Rightarrow (iii) As an immediate consequence of Lemma 2.13 one verifies that the map $\Theta : \pi_0(\text{End}(A \otimes \mathbb{K})) \rightarrow K_0(A)$, $\Theta[\alpha] = [\alpha(1 \otimes e)]$ is multiplicative, i.e. $\Theta[\alpha \circ \beta] = \Theta[\alpha]\Theta[\beta]$. Let $q := \alpha^{-1}(1 \otimes e)$. Then $[p][q] = \Theta[\alpha \circ \alpha^{-1}] = \Theta[\text{id}] = [1]$.

(iii) \Rightarrow (i) By assumption there is a full projection $q \in A \otimes K$ such that $[p][q] = [1]$ in $K_0(A)$. By Lemma 2.10 there are $\varphi, \psi \in \text{End}(A \otimes \mathbb{K})$ such that $\varphi(1 \otimes e) = p$ and $\psi(1 \otimes e) = q$. Since $[p][q] = [1]$ in $K_0(A)$, it follows that $[\varphi \circ \psi] = [\psi \circ \varphi] = [\text{id}_{A \otimes \mathbb{K}}] \in [A \otimes \mathbb{K}, A \otimes \mathbb{K}]$. Therefore $\varphi \circ \psi \approx_u \text{id}_{A \otimes \mathbb{K}} \approx_u \psi \circ \varphi$ by Proposition 2.12. By [38, Cor.2.3.4] it follows that there is an automorphism $\varphi_0 \in \text{Aut}(A \otimes \mathbb{K})$ such that $\varphi_0 \approx_u \varphi$. Set $p_0 = \varphi_0(1_A \otimes e)$. The map φ_0 induces a $*$ -isomorphism $A \cong (1_A \otimes e)(A \otimes \mathbb{K})(1_A \otimes e) \rightarrow p_0(A \otimes \mathbb{K})p_0$. We conclude that $A \cong p(A \otimes \mathbb{K})p$ since p_0 is unitarily equivalent to p . \square

If A is a separable unital C^* -algebra, Brown, Green and Rieffel [10] showed that the Picard group $\text{Pic}(A)$ is isomorphic to the outer automorphism group of $A \otimes \mathbb{K}$, i.e. $\text{Pic}(A) \cong \text{Out}(A \otimes \mathbb{K}) = \text{Aut}(A \otimes \mathbb{K})/\text{Inn}(A \otimes \mathbb{K})$. One can view $\text{Out}(A)$ as a subgroup of $\text{Pic}(A)$. Kodaka [26] has shown that the coset space $\text{Pic}(A)/\text{Out}(A)$ is in bijection with the Murray-von Neumann equivalence classes of full projections $p \in A \otimes \mathbb{K}$ such that $p(A \otimes \mathbb{K})p \cong A$. From Lemmas 2.13 and 2.14 we see that if A is strongly self-absorbing, then $\text{Out}(A)$ is a normal subgroup of $\text{Pic}(A)$ and we have:

Corollary 2.15. *If A is strongly self-absorbing, then there is an exact sequence of groups*

$$1 \rightarrow \text{Out}(A) \rightarrow \text{Pic}(A) \rightarrow K_0(A)_+^\times \rightarrow 1.$$

If moreover A is stably finite, then its normalized trace induces a homomorphism of multiplicative groups from $K_0(A)_+^\times$ onto the fundamental group $\mathcal{F}(A)$ of A defined in [35].

Lemma 2.16. *Let A be a strongly self-absorbing C^* -algebra. The sequence $\text{Aut}_{1 \otimes e}(A \otimes \mathbb{K}) \rightarrow \text{Aut}(A \otimes \mathbb{K}) \rightarrow \text{Pr}(A \otimes \mathbb{K})^\times$ where the first map is the inclusion and the second sends α to $\alpha(1 \otimes e)$ is a locally trivial fiber bundle over a paracompact base space and therefore it is a Hurewicz fibration.*

Proof. Lemma 2.14 shows that the map to the base space is surjective. With this remark, the proof is entirely similar the proof of Lemma 2.8. \square

Corollary 2.17. *Let A be a strongly self-absorbing C^* -algebra. Then $\text{Aut}(A \otimes \mathbb{K})$ has the homotopy type of a CW complex, which is homotopic to $K_0(A)_+^\times \times BU(A)$.*

Proof. $\text{Aut}(A \otimes \mathbb{K})^\times$ is the coproduct of its path components, all of which are homeomorphic to $\text{Aut}_0(A \otimes \mathbb{K})$. By Theorem 2.5 and Lemma 2.8, $\text{Aut}_0(A \otimes \mathbb{K})$ is homotopy equivalent to $\mathcal{P}r_0(A \otimes \mathbb{K})$. By Lemma 2.14, $\pi_0(\mathcal{P}r(A \otimes \mathbb{K})^\times) \cong K_0(A)_+^\times$. Thus, using Corollary 2.9 we have

$$\text{Aut}(A \otimes \mathbb{K}) \simeq \pi_0(\mathcal{P}r(A \otimes \mathbb{K})^\times) \times \mathcal{P}r_0(A \otimes \mathbb{K}) \simeq K_0(A)_+^\times \times BU(A). \quad \square$$

In the case $A = \mathbb{C}$ this reproves the well-known fact that $\text{Aut}(\mathbb{K}) \simeq BU(1) \simeq K(\mathbb{Z}, 2)$ and hence the only non vanishing homotopy group of $\text{Aut}(\mathbb{K})$ is $\pi_2(\text{Aut}(\mathbb{K})) \cong \pi_2(BU(1)) \cong \pi_1(U(1)) \cong \mathbb{Z}$. At the same time, for $A \neq \mathbb{C}$, we obtain the following.

Theorem 2.18. *Let $A \neq \mathbb{C}$ be a strongly self-absorbing C^* -algebra. Then there are isomorphisms of groups*

$$\pi_i(\text{Aut}(A \otimes \mathbb{K})) = \begin{cases} K_0(A)_+^\times & \text{if } i = 0 \\ K_i(A) & \text{if } i \geq 1. \end{cases}$$

Proof. We have seen in the proof of Corollary 2.17 that $\pi_0(\text{Aut}(A \otimes \mathbb{K})) \cong K_0(A)_+^\times$. If $i \geq 1$, then by Corollary 2.17, $\pi_i(\text{Aut}(A \otimes \mathbb{K})) \cong \pi_i(BU(A)) \cong \pi_{i-1}(U(A))$. On the other hand, since A is \mathcal{Z} -stable, we have that $\pi_{i-1}(U(A)) \cong K_i(A)$ by [25, Thm.3]. \square

Corollary 2.19. *Let A be a strongly self-absorbing C^* -algebra. There is an exact sequence of topological groups $1 \rightarrow \text{Aut}_0(A \otimes \mathbb{K}) \rightarrow \text{Aut}(A \otimes \mathbb{K}) \rightarrow K_0(A)_+^\times \rightarrow 1$.*

Remark 2.20. The exact sequence $1 \rightarrow \text{Aut}_0(\mathcal{O}_\infty \otimes \mathbb{K}) \rightarrow \text{Aut}(\mathcal{O}_\infty \otimes \mathbb{K}) \rightarrow \mathbb{Z}/2 \rightarrow 1$ is split, since by [6] there is an order-two automorphism α of $\mathcal{O}_\infty \otimes \mathbb{K}$ such that $\alpha_* = -1$ on $K_0(\mathcal{O}_\infty)$.

Corollary 2.21. *Let $A \neq \mathbb{C}$ be a strongly self-absorbing C^* -algebra. The natural map $\text{Aut}_0(A \otimes \mathbb{K}) \rightarrow \text{Aut}_0(\mathcal{O}_\infty \otimes A \otimes \mathbb{K})$ is a homotopy equivalence.*

Proof. $\text{Aut}_0(A \otimes \mathbb{K}) \rightarrow \text{Aut}_0(\mathcal{O}_\infty \otimes A \otimes \mathbb{K})$ is given by $\alpha \mapsto \text{id}_{\mathcal{O}_\infty} \otimes \alpha$. Both spaces have the homotopy type of a CW-complex by Corollary 2.9 and they are weakly homotopic by Thm. 2.18. \square

Theorem 2.22. *Let $A \neq \mathbb{C}$ be a strongly self-absorbing C^* -algebra and let X be a compact metrizable space. The map $\Theta: [X, \text{End}(A \otimes \mathbb{K})] \rightarrow K_0(C(X) \otimes A)_+$ given by $\Theta([\alpha]) = [\alpha(1 \otimes e)]$ is an isomorphism of commutative semirings. Θ restricts to a group isomorphism $[X, \text{Aut}(A \otimes \mathbb{K})] \rightarrow K_0(C(X) \otimes A)_+^\times$. If X is connected, then $K_0(C(X) \otimes A)_+^\times \cong K_0(A)_+^\times \oplus K_0(C_0(X \setminus x_0) \otimes A)$. If A is purely infinite, then $K_0(C(X) \otimes A)_+ = K_0(C(X) \otimes A)$ and Θ is an isomorphism of rings.*

Proof. Let $\psi: A \otimes \mathbb{K} \rightarrow (A \otimes \mathbb{K})^{\otimes 2}$ and l be as in Lemma 2.13. The additivity of Θ is easily verified. Let $\alpha, \beta \in C(X, \text{End}(A \otimes \mathbb{K}))$, then by Lemma 2.13:

$$\begin{aligned} [(\alpha \circ \beta)(1 \otimes e)] &= [(\alpha * \beta)(1 \otimes e)] = [\psi^{-1} \circ (\alpha \otimes \beta) \circ \psi(1 \otimes e)] \\ &= [\psi^{-1}(\alpha(1 \otimes e) \otimes \beta(1 \otimes e))] = [\alpha(1 \otimes e)] \cdot [\beta(1 \otimes e)], \end{aligned}$$

which shows that $\Theta: [X, \text{End}(A \otimes \mathbb{K})] \rightarrow K_0(C(X) \otimes A)_+$ is a homomorphism of semirings. Let $p \in C(X) \otimes A \otimes \mathbb{K}$ be a full projection. Then, $p(C(X) \otimes A \otimes \mathbb{K})p$ is A -absorbing by [49, Cor.3.1]. It follows that Θ is surjective by Lemma 2.10. For injectivity we apply Proposition 2.12(i).

Next we show that the image of the restriction of Θ to $[X, \text{Aut}(A \otimes \mathbb{K})]$ coincides with $K_0(C(X) \otimes \mathbb{K})_+^\times$. Let $p \in \mathcal{P}r(C(X) \otimes A \otimes \mathbb{K})^\times$. By assumption, there is $q \in \mathcal{P}r(C(X) \otimes A \otimes \mathbb{K})^\times$ such that $[p][q] = 1$ in the ring $K_0(C(X) \otimes A)$. By Lemma 2.10 there are $\varphi, \psi \in \text{Hom}(A \otimes \mathbb{K}, C(X) \otimes A \otimes \mathbb{K})$ such that $\varphi(1 \otimes e) = p$ and $\psi(1 \otimes e) = q$. Let $\tilde{\varphi}, \tilde{\psi} \in \text{End}_{C(X)}(C(X) \otimes A \otimes \mathbb{K})$ be the unique $C(X)$ -linear extensions of φ and ψ . Note that if $\iota : A \otimes \mathbb{K} \rightarrow C(X) \otimes A \otimes \mathbb{K}$ is the inclusion $\iota(a) = 1_{C(X)} \otimes a$, then $\tilde{\iota} = \text{id}_{C(X) \otimes A \otimes \mathbb{K}}$. Since $[p][q] = [1]$ in $K_0(C(X) \otimes A)$, it follows that $[\tilde{\varphi} \circ \psi] = [\tilde{\psi} \circ \varphi] = [\iota] \in [A \otimes \mathbb{K}, C(X) \otimes A \otimes \mathbb{K}]$. By Proposition 2.12(ii) it follows that $\tilde{\varphi} \circ \psi \approx_u \iota \approx_u \tilde{\psi} \circ \varphi$. This clearly implies that $\tilde{\varphi} \circ \tilde{\psi} \approx_u \text{id}_{C(X) \otimes A \otimes \mathbb{K}} \approx_u \tilde{\psi} \circ \tilde{\varphi}$. By [38, Cor.2.3.4] it follows that there is an automorphism $\alpha \in \text{Aut}_{C(X)}(C(X) \otimes A \otimes \mathbb{K})$ such that $\alpha \approx_u \tilde{\varphi}$. In particular we have that $[\alpha(1 \otimes e)] = [\tilde{\varphi}(1 \otimes e)] = [p]$.

It remains to verify the isomorphism $K_0(C(X) \otimes A)_+^\times \cong K_0(A)_+^\times \oplus K_0(C_0(X \setminus x_0) \otimes A)$. Evaluation at x_0 induces a split exact sequence $0 \rightarrow K_0(C_0(X \setminus x_0) \otimes A) \rightarrow K_0(C(X) \otimes A) \rightarrow K_0(A) \rightarrow 0$. Arguing as in the proof of [13, Prop.5.6], one verifies that $K_0(C_0(X \setminus x_0) \otimes A)$ is a nil-ideal of the ring $K_0(C(X) \otimes A)$. Thus an element $\sigma \in K_0(C(X) \otimes A)$ is invertible if and only if its restriction $\sigma_{x_0} \in K_0(A)$ is invertible. Consequently $K_0(C(X) \otimes A)^\times \cong K_0(A)^\times \oplus K_0(C_0(X \setminus x_0) \otimes A)$. It remains to verify that an element $\sigma \in K_0(C(X) \otimes A)$ is positive if $\sigma_{x_0} \in K_0(A)_+ \setminus \{0\}$. It suffices to consider the case when A is stably finite. Let τ denote the unique trace state of A . Its extension to a trace state on $A \otimes M_n(\mathbb{C})$ is denoted again by τ . Then any continuous trace η on $C(X) \otimes A \otimes M_n(\mathbb{C})$ is of the form $\eta(f) = \int_X \tau(f) d\mu$ for some finite Borel measure μ on X . Write $\sigma = [p] - [q]$ where $p, q \in \mathcal{P}r(C(X) \otimes A \otimes M_n(\mathbb{C}))$ are full projections. Let r be a nonzero projection in $A \otimes M_n(\mathbb{C})$ such that $[p(x_0)] - [q(x_0)] = [r]$. Since X is connected it follows that $[p(x)] - [q(x)] = [r] \in K_0(A)$ for all $x \in X$. From this we see that any point $x \in X$ has a closed neighborhood V such that $[p_V] - [q_V] = [r] \in K_0(C(V) \otimes A)$. Since $\tau(r) > 0$ it follows immediately that $\eta(p) > \eta(q)$ for all nonzero finite traces η on $A \otimes M_n(\mathbb{C})$. We apply Corollaries 4.9 and 4.10 of [39] to conclude that $[p] - [q] \in K_0(C(X) \otimes A)_+$. \square

Corollary 2.23. *Let X be a compact connected metrizable space. Then there are isomorphism of multiplicative groups*

$$\begin{aligned} [X, \text{Aut}(\mathcal{Z} \otimes \mathbb{K})] &\cong K^0(X)_+^\times = 1 + \tilde{K}^0(X), \\ [X, \text{Aut}(\mathcal{O}_\infty \otimes \mathbb{K})] &\cong K^0(X)^\times = \pm 1 + \tilde{K}^0(X). \end{aligned}$$

2.5. The topological group $\text{Aut}(A \otimes \mathbb{K})$ is well-pointed. Since we would like to apply the nerve construction to deloop the topological group $\text{Aut}(A \otimes \mathbb{K})$, we will need to show that $\text{Aut}(A \otimes \mathbb{K})$ is well-pointed. This notion is defined as follows:

Definition 2.24. Let X be a topological space, $A \subset X$ a closed subspace. The pair (X, A) is called a *neighborhood deformation retract* (or *NDR-pair* for short) if there is a map $u : X \rightarrow I = [0, 1]$ such that $u^{-1}(0) = A$ and a homotopy $H : X \times I \rightarrow X$ such that $H(x, 0) = x$ for all $x \in X$, $H(a, t) = a$ for $a \in A$ and $t \in I$ and $H(x, 1) \in A$ if $u(x) < 1$. A pointed topological space X with basepoint $x_0 \in X$ is said to have a *non-degenerate basepoint* or to be *well-pointed* if the pair (X, x_0) is an NDR-pair.

Recall that a neighborhood V of x_0 deformation retracts to x_0 if there is a continuous map $h : V \times I \rightarrow V$ such that $h(x, 0) = v$, $h(x_0, t) = x_0$ and $h(x, 1) = x_0$ for all $x \in V$ and $t \in I$. The following lemma is contained in [46, Thm.2].

Lemma 2.25. *Let (X, x_0) be a pointed topological space together with a continuous map $v : X \rightarrow I$ such that $x_0 = v^{-1}(0)$ and $V = \{x \in X : v(x) < 1\}$ deformation retracts to x_0 . Then (X, x_0) is an NDR-pair.*

Proposition 2.26. *Let A be a strongly self-absorbing C^* -algebra. Then the topological group $\text{Aut}(A \otimes \mathbb{K})$ is well-pointed.*

Proof. let $e \in \mathbb{K}$ be a rank-1 projection and set $p_0 = 1 \otimes e$. Let $U = \{p \in \mathcal{P}r(A \otimes \mathbb{K}) \mid \|p - p_0\| < 1/2\}$. If $\pi : \text{Aut}_0(A \otimes \mathbb{K}) \rightarrow \mathcal{P}r_0(A \otimes \mathbb{K})$ denotes the map $\pi \mapsto \pi(p_0)$, we will show that $\pi^{-1}(U)$ deformation retracts to $\text{id}_{A \otimes \mathbb{K}} \in \text{Aut}_0(A \otimes \mathbb{K})$. As we have seen in the proof of Lemma 2.8, the principal bundle $\text{Aut}_0(A \otimes \mathbb{K}) \rightarrow \mathcal{P}r_0(A \otimes \mathbb{K})$ trivializes over U , i.e. there exists a homeomorphism $\pi^{-1}(U) \rightarrow U \times \text{Aut}_{p_0}(A \otimes \mathbb{K})$ sending $\text{id}_{A \otimes \mathbb{K}}$ to $(p_0, \text{id}_{A \otimes \mathbb{K}})$. Thus, it suffices to show that the right hand side retracts. Let χ be the characteristic function of $(1/2, 1]$. Then $h(p, t) = \chi((1-t)p + tp_0)$ is a deformation retraction of U into p_0 . This is well-defined since $1/2$ is not in the spectrum of $a = (1-t)p + tp_0$ as seen from the estimate $\|(1-2a) - (1-2p_0)\| < 1$. We have shown in Theorem 2.5 that $\text{Aut}_{p_0}(A \otimes \mathbb{K})$ deformation retracts to $\text{id}_{A \otimes \mathbb{K}}$. Combining these homotopies we end up with a deformation retraction of $\pi^{-1}(U)$ into $\text{id}_{A \otimes \mathbb{K}}$. Let d be a metric for $\text{Aut}(A \otimes \mathbb{K})$. Then $v : \text{Aut}(A \otimes \mathbb{K}) \rightarrow [0, 1]$, $v(\alpha) = \max\{\min\{d(\alpha, \text{id}_{A \otimes \mathbb{K}}), 1/2\}, \min\{1, 2\|\alpha(p_0) - p_0\|\}\}$ and $V := \pi^{-1}(U)$ satisfy the conditions of Lemma 2.25 relative to the basepoint $\text{id}_{A \otimes \mathbb{K}}$. \square

3. THE INFINITE LOOP SPACE STRUCTURE OF $B\text{Aut}(A \otimes \mathbb{K})$

3.1. Permutative categories and infinite loop spaces. We will show that $B\text{Aut}(A \otimes \mathbb{K})$ is an infinite loop space in the sense of the following definition [1].

Definition 3.1. A topological space $E = E_0$ is called an *infinite loop space*, if there exists a sequence of spaces E_i , $i \in \mathbb{N}$, such that $E_i \simeq \Omega E_{i+1}$ for all $i \in \mathbb{N}_0$ (\simeq denotes homotopy equivalence).

The importance of these spaces lies in the fact, that they represent generalized cohomology theories, i.e. for a CW-complex X , the homotopy classes of maps $E^i(X) := [X, E_i]$ are abelian groups and the functor $X \mapsto E^\bullet(X)$ is a cohomology theory. The sequence of spaces E_i forms a *connective Ω -spectrum*. There is a well-developed theory to detect whether a space belongs to this class [29, 44]. One of the main sources for infinite loop spaces are classifying spaces of topological strict symmetric monoidal categories, called *permutative categories* in [31].

A *topological category* has a space of objects, a space of morphisms and continuous source, target and identity maps. Such a category \mathcal{C} carries a *strict monoidal structure* if it comes equipped with a functor $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ that satisfies the analogues of associativity and unitality known for monoids. The strictness refers to the fact that these hold on the nose, not only up to natural transformations. \mathcal{C} is called *symmetric* if it comes equipped with a natural transformation $c : \otimes \circ \tau \rightarrow \otimes$, where $\tau : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C} \times \mathcal{C}$ is switching the factors. This should behave like a permutation on

n -fold tensor products. We will assume all permutative categories to be *well-pointed* in the sense that the map $\text{obj}(\mathcal{C}) \rightarrow \text{mor}(\mathcal{C})$, $x \rightarrow \text{id}_x$ is a cofibration. For a precise definition, we refer the reader to [31, Def.1].

Any topological category \mathcal{C} can be turned into a simplicial space $N_\bullet \mathcal{C}$ via the nerve construction. Let $N_0 \mathcal{C} = \text{obj}(\mathcal{C})$, $N_1 \mathcal{C} = \text{mor}(\mathcal{C})$ and

$$N_k \mathcal{C} = \{(f_1, \dots, f_k) \in \text{mor}(\mathcal{C}) \times \dots \times \text{mor}(\mathcal{C}) \mid s(f_i) = t(f_{i+1})\}.$$

The face maps $d_i^k: N_k \mathcal{C} \rightarrow N_{k-1} \mathcal{C}$ and degeneracies $s_i^k: N_k \mathcal{C} \rightarrow N_{k+1} \mathcal{C}$ are induced by composition of successive maps and insertion of identities respectively. The geometric realization of a simplicial space X_\bullet is defined by

$$|X_\bullet| = \left(\prod_{k=0} X_k \times \Delta_k \right) / \sim$$

where $\Delta_k \subset \mathbb{R}^{k+1}$ denotes the standard k -simplex and the equivalence relation is generated by $(d_i^k x, u) \sim (x, \partial_i^{k-1} u)$ and $(s_i^l y, v) \sim (y, \sigma_i^{l+1} v)$ for $x \in X_k$, $u \in \Delta_{k-1}$, $y \in X_l$, $v \in \Delta_{l+1}$, where δ_i and σ_i are the coface and codegeneracy maps on the standard simplex. For details about this construction we refer the reader to [29, sec.11].

The space $|N_\bullet \mathcal{C}|$ associated to a category \mathcal{C} is called the *classifying space* of \mathcal{C} . Having a monoidal structure on \mathcal{C} yields the following.

Lemma 3.2. *Let \mathcal{C} be a strict monoidal topological category. Then $|N_\bullet \mathcal{C}|$ is a topological monoid.*

Proof. The nerve construction N_\bullet preserves products in the sense that the projection functors $\pi_i: \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ induce a levelwise homeomorphism $N_\bullet(\mathcal{C} \times \mathcal{C}) \rightarrow N_\bullet \mathcal{C} \times N_\bullet \mathcal{C}$. Therefore $N_\bullet \mathcal{C}$ is a simplicial topological monoid and the lemma follows from [29, Cor.11.7]. \square

A permutative category \mathcal{C} provides an input for infinite loop space machines [31, Def.2]. Due to the above lemma, there is a classifying space $B|N_\bullet \mathcal{C}|$. The following has been proven by Segal [44] and May [30, Thm.4.10]

Theorem 3.3. *Let \mathcal{C} be a permutative category. Then $\Omega B|N_\bullet \mathcal{C}|$ is an infinite loop space. Moreover, if $\pi_0(|N_\bullet \mathcal{C}|)$ is a group, then the map $|N_\bullet \mathcal{C}| \rightarrow \Omega B|N_\bullet \mathcal{C}|$ induced by the inclusion of the 1-skeleton $S^1 \times |N_\bullet \mathcal{C}| \rightarrow B|N_\bullet \mathcal{C}|$ is a homotopy equivalence.*

We define a *non-unital permutative category* to be a category that satisfies all the axioms of a permutative category, but does not have a unit object. We can turn this into an honest permutative category \mathcal{C}^+ , called the *unitization* of \mathcal{C} , via $\text{obj}(\mathcal{C}^+) = \text{obj}(\mathcal{C}) \amalg \{p_0\}$ and $\text{mor}(\mathcal{C}^+) = \text{mor}(\mathcal{C}) \amalg \{\text{id}_{p_0}\}$. The new source and target maps send id_{p_0} to p_0 and are the source and target maps of \mathcal{C} on the rest, similarly for the identity map.

The monoidal structure can be extended to \mathcal{C}^+ via $p_0 \otimes x = x$ and $\text{id}_{p_0} \otimes f = f$ for $x \in \text{obj}(\mathcal{C})$ and $f \in \text{mor}(\mathcal{C})$. Likewise, the symmetry c can be extended to \mathcal{C}^+ via $c_{p_0, x} = \text{id}_x = c_{x, p_0}$. It is straightforward to check that \mathcal{C}^+ is indeed a permutative category. We will use the following lemma, which can be found in a slightly different context in [47, Lemma A.1].

Lemma 3.4. *Let \mathcal{C} be a non-unital permutative category and assume that $|N_\bullet \mathcal{C}|$ is a homotopy unital grouplike H -space with respect to the operation induced by the tensor product on \mathcal{C} . Let \mathcal{C}^+ be the unitization of \mathcal{C} . Then the inclusion functor $\mathcal{C} \rightarrow \mathcal{C}^+$ induces a weak homotopy equivalence*

$$|N_\bullet \mathcal{C}| \rightarrow \Omega B(|N_\bullet \mathcal{C}^+|) .$$

In particular, if $\text{obj}(\mathcal{C})$ and $\text{mor}(\mathcal{C})$ have the homotopy type of CW complexes, then $|N_\bullet \mathcal{C}|$ is an infinite loop space.

Proof. Let $G = \pi_0(|N_\bullet \mathcal{C}|)$. Then we have $G' = \pi_0(|N_\bullet \mathcal{C}^+|) = G \cup \{x\}$ with monoid structure given by $g \cdot x = x \cdot g = g$, $x \cdot x = x$ and the usual group multiplication in G . Thus, we may think of the element x as a “second unit element” in G' . The group completion of this monoid consists of equivalence classes of pairs $(g, h) \in G'$, where

$$(g, h) \sim (g', h') \quad \text{if and only if} \quad \exists y \in G' : gh'y = g'hy .$$

In particular, we have $(x, e) \sim (e, e)$, where $e \in G \subset G'$ is the ordinary unit element. Therefore, the inclusion map $G \rightarrow G'$ induces an isomorphism on the group completions, with the inverse induced by the map from $G' \rightarrow G$ that sends x to e .

Since both sides are H -spaces and therefore simple (i.e. π_1 acts trivially on all higher homotopy groups), the result will follow from Whitehead’s theorem, if we can prove that $H_*(|N_\bullet \mathcal{C}|) \rightarrow H_*(\Omega B|N_\bullet \mathcal{C}^+|)$ is an isomorphism. Note that both sides are actually commutative rings due to the multiplicative structures on both spaces and the symmetry on \mathcal{C} . By the Segal-McDuff group completion theorem [33, Prop.1] the inclusion $|N_\bullet \mathcal{C}^+| \rightarrow \Omega B|N_\bullet \mathcal{C}^+|$ yields a ring isomorphism

$$H_*(|N_\bullet \mathcal{C}^+|)[\pi_0^{-1}] \rightarrow H_*(\Omega B|N_\bullet \mathcal{C}^+|) ,$$

where $\pi_0 \subset H_0(|N_\bullet \mathcal{C}^+|) \subset H_*(|N_\bullet \mathcal{C}^+|)$ is the multiplicative subset of the path components in the homology ring. A similar line of thought as the one given above shows that the map into the localization

$$H_*(|N_\bullet \mathcal{C}|) \rightarrow H_*(|N_\bullet \mathcal{C}^+|)[\pi_0^{-1}]$$

is a ring isomorphism. The last statement is a consequence of [44, Prop.A.1 (i), (iv)] together with the fact that $N_\bullet \mathcal{C}$ is good for well-pointed categories \mathcal{C} . \square

3.2. The tensor product of $A \otimes \mathbb{K}$ -bundles. Let A be a strongly self-absorbing C^* -algebra, X be a topological space and let P_1 and P_2 be principal $\text{Aut}(A \otimes \mathbb{K})$ -bundles over X . Fix an isomorphism $\psi: A \otimes \mathbb{K} \rightarrow (A \otimes \mathbb{K}) \otimes (A \otimes \mathbb{K})$. This choice induces a tensor product operation on principal $\text{Aut}(A \otimes \mathbb{K})$ -bundles in the following way. Note that $P_1 \times_X P_2 \rightarrow X$ is a principal $\text{Aut}(A \otimes \mathbb{K}) \times \text{Aut}(A \otimes \mathbb{K})$ -bundle and ψ induces a group homomorphism

$$\text{Ad}_{\psi^{-1}}: \text{Aut}(A \otimes \mathbb{K}) \times \text{Aut}(A \otimes \mathbb{K}) \rightarrow \text{Aut}(A \otimes \mathbb{K}) \quad ; \quad (\alpha, \beta) \mapsto \psi^{-1} \circ (\alpha \otimes \beta) \circ \psi .$$

Now let

$$P_1 \otimes_\psi P_2 := (P_1 \times_X P_2) \times_{\text{Ad}_{\psi^{-1}}} \text{Aut}(A \otimes \mathbb{K}) = ((P_1 \times_X P_2) \times \text{Aut}(A \otimes \mathbb{K})) / \sim$$

where the equivalence relation is $(p_1 \alpha, p_2 \beta, \gamma) \sim (p_1, p_2, \text{Ad}_{\psi^{-1}}(\alpha, \beta) \gamma)$ for all $(p_1, p_2) \in P_1 \times_X P_2$ and $\alpha, \beta, \gamma \in \text{Aut}(A \otimes \mathbb{K})$. This is a delooped version of the operation $*$ from Lemma 2.13.

Due to the choice of ψ , which was arbitrary, \otimes_ψ can not be associative. We will show, however, that – just like $*$ – it is *homotopy associative* and also *homotopy unital*.

To obtain a model for the classifying space $B\text{Aut}(A \otimes \mathbb{K})$, let \mathcal{B} be the topological category, which has as its object space just a single point and the group $\text{Aut}(A \otimes \mathbb{K})$ as its morphism space. Since we have shown that $\text{Aut}(A \otimes \mathbb{K})$ is well-pointed (see Proposition 2.26), [32, Prop.7.5 and Thm.8.2] implies that the geometric realization $|N_\bullet \mathcal{B}|$ has in fact the homotopy type of a classifying space for principal $\text{Aut}(A \otimes \mathbb{K})$ -bundles, i.e.

$$B\text{Aut}(A \otimes \mathbb{K}) = |N\mathcal{B}_\bullet|.$$

Choosing an isomorphism $\psi: A \otimes \mathbb{K} \rightarrow (A \otimes \mathbb{K}) \otimes (A \otimes \mathbb{K})$, we can define a functor $\otimes_\psi: \mathcal{B} \times \mathcal{B} \rightarrow \mathcal{B}$ just as above, which acts on morphisms $\alpha, \beta \in \text{Aut}(A \otimes \mathbb{K})$ by

$$(\alpha, \beta) \mapsto \alpha \otimes_\psi \beta := \text{Ad}_{\psi^{-1}}(\alpha, \beta).$$

This is in fact functorial since composition is well-behaved with respect to the tensor product in the following way

$$\begin{aligned} (\alpha \circ \alpha') \otimes_\psi (\beta \circ \beta') &= \psi^{-1} \circ (\alpha \circ \alpha') \otimes (\beta \circ \beta') \circ \psi \\ &= \psi^{-1} \circ (\alpha \otimes \beta) \circ \psi \circ \psi^{-1} \circ (\alpha' \otimes \beta') \circ \psi = (\alpha \otimes_\psi \beta) \circ (\alpha' \otimes_\psi \beta'). \end{aligned}$$

The functor induces a multiplication map on the geometric realization

$$\mu_\psi: B\text{Aut}(A \otimes \mathbb{K}) \times B\text{Aut}(A \otimes \mathbb{K}) \rightarrow B\text{Aut}(A \otimes \mathbb{K}).$$

Observe that a path connecting two isomorphisms $\psi, \psi' \in \text{Iso}(\text{Aut}(A \otimes \mathbb{K}), \text{Aut}(A \otimes \mathbb{K})^{\otimes 2})$ induces a homotopy of functors $\mathcal{B} \times \mathcal{B} \times I \rightarrow \mathcal{B}$, where I here is the category, which has $[0, 1]$ as its object space and only identities as morphisms. After geometric realization this in turn yields a homotopy between μ_ψ and $\mu_{\psi'}$ (observe that $|I| \cong [0, 1]$).

Lemma 3.5. *Let A be a strongly self-absorbing C^* -algebra and let \mathcal{B} be the category defined above. Let $\psi: A \otimes \mathbb{K} \rightarrow (A \otimes \mathbb{K}) \otimes (A \otimes \mathbb{K})$ be an isomorphism, then \otimes_ψ defines an H -space structure on $B\text{Aut}(A \otimes \mathbb{K})$, which has the basepoint of $B\text{Aut}(A \otimes \mathbb{K})$ as a homotopy unit. Different choices of ψ yield homotopy equivalent H -space structures.*

Proof. The proof of this statement is very similar to the one of Lemma 2.13, but we have to take care that the homotopies we use run through *functors* on \mathcal{B} . Let $\hat{\psi}, \hat{l}$ and \hat{r} be just as in Theorem 2.5 and consider $\psi = \hat{\psi}(\frac{1}{2})$ first. By Theorem 2.5, there is a path between $(\psi \otimes \text{id}_{A \otimes \mathbb{K}}) \circ \psi$ and $(\text{id}_{A \otimes \mathbb{K}} \otimes \psi) \circ \psi$, since both these morphisms map $1 \otimes e$ to $(1 \otimes e)^{\otimes 3}$. This proves the homotopy associativity in this case.

To prove that the basepoint provides a homotopy unit we have to show that the two functors $\alpha \mapsto \alpha \otimes_\psi \text{id}_{A \otimes \mathbb{K}}$ and $\alpha \mapsto \text{id}_{A \otimes \mathbb{K}} \otimes_\psi \alpha$ are both homotopic to the identity functor. The argument for this is the same as in the proof of Lemma 2.13.

Now let $\psi: A \otimes \mathbb{K} \rightarrow (A \otimes \mathbb{K}) \otimes (A \otimes \mathbb{K})$ be an arbitrary isomorphism. As in Lemma 2.13 we have $\psi = \hat{\psi} \circ \kappa$ for some automorphism $\kappa \in \text{Aut}(A \otimes \mathbb{K})$. If we denote homotopic functors by \sim , we have

$$\begin{aligned} \alpha \otimes_{\psi} \beta &= \kappa^{-1} \circ (\alpha \otimes_{\hat{\psi}} \beta) \circ \kappa \sim (\kappa \otimes_{\hat{\psi}} \text{id}_{A \otimes \mathbb{K}})^{-1} \circ (\text{id}_{A \otimes \mathbb{K}} \otimes_{\hat{\psi}} (\alpha \otimes_{\hat{\psi}} \beta)) \circ (\kappa \otimes_{\hat{\psi}} \text{id}_{A \otimes \mathbb{K}}) \\ &= \text{id}_{A \otimes \mathbb{K}} \otimes_{\hat{\psi}} (\alpha \otimes_{\hat{\psi}} \beta) \sim (\alpha \otimes_{\hat{\psi}} \beta) . \end{aligned}$$

Note that every stage of this homotopy provides functors $\mathcal{B} \times \mathcal{B} \rightarrow \mathcal{B}$. Geometrically realizing this homotopy we see that different choices of ψ yield the same H -space structure up to homotopy. \square

To apply the infinite loop space machine, we need an honest permutative category encoding the operation \otimes_{ψ} . Let \mathcal{B}_{\otimes} be the category, which has \mathbb{N} as its object space (where $n \in \mathbb{N}$ should be thought of as $A^{\otimes n}$). The morphisms between n and m are given by $\text{Iso}((A \otimes \mathbb{K})^{\otimes n}, (A \otimes \mathbb{K})^{\otimes m})$ equipped with the point-norm topology. In particular, \mathcal{B}_{\otimes} is a groupoid and we have $\text{hom}(1, 1) = \text{Aut}(A \otimes \mathbb{K})$.

The tensor product induces a *strict* monoidal structure on \mathcal{B}_{\otimes} with $n \otimes m = n + m$ and the ordinary tensor product on morphisms. Likewise, we have a symmetry $c_{n,m}$ on \mathcal{B}_{\otimes} , where $c_{n,m} \in \text{Aut}((A \otimes \mathbb{K})^{\otimes(n+m)})$ is the automorphism $(A \otimes \mathbb{K})^{\otimes n} \otimes (A \otimes \mathbb{K})^{\otimes m} \rightarrow (A \otimes \mathbb{K})^{\otimes m} \otimes (A \otimes \mathbb{K})^{\otimes n}$ switching the two factors. This turns \mathcal{B}_{\otimes} into a non-unital permutative category and as above we denote its unitization by \mathcal{B}_{\otimes}^+ , which is still a groupoid. Let

$$BAut(A \otimes \mathbb{K})_{\otimes} = |N_{\bullet} \mathcal{B}_{\otimes}| \quad \text{and} \quad BAut(A \otimes \mathbb{K})_{\otimes}^+ = |N_{\bullet} \mathcal{B}_{\otimes}^+| .$$

In the following lemma we use the fact that an equivalence of categories induces a homotopy equivalence of classifying spaces. This is a direct corollary of [43, Prop.2.1].

Lemma 3.6. *The inclusion functor $J: \mathcal{B} \rightarrow \mathcal{B}_{\otimes}$ induces an equivalence of categories with continuous inverse functor. Given an isomorphism $\psi: A \otimes \mathbb{K} \rightarrow (A \otimes \mathbb{K})^{\otimes 2}$, the diagram*

$$(2) \quad \begin{array}{ccc} \mathcal{B} \times \mathcal{B} & \xrightarrow{\otimes_{\psi}} & \mathcal{B} \\ J \times J \downarrow & & \downarrow J \\ \mathcal{B}_{\otimes} \times \mathcal{B}_{\otimes} & \xrightarrow{\otimes} & \mathcal{B}_{\otimes} \end{array}$$

commutes up to a natural transformation. In particular we have a homotopy equivalence, $BAut(A \otimes \mathbb{K}) \simeq BAut(A \otimes \mathbb{K})_{\otimes}$. Moreover, we have $BAut(A \otimes \mathbb{K})_{\otimes}^+ \simeq BAut(A \otimes \mathbb{K})_{\otimes} \cup \{pt\}$.

Proof. The last statement is clear from the definition of the nerve and the geometric realization. To see the first claim, choose isomorphisms $\psi_n: A \otimes \mathbb{K} \rightarrow (A \otimes \mathbb{K})^{\otimes n}$ for every $n \in \mathbb{N}$ with $\psi_n(1 \otimes e) = (1 \otimes e)^{\otimes n}$ and $\psi_1 = \text{id}_{A \otimes \mathbb{K}}$. Now define a functor $R: \mathcal{B}_{\otimes} \rightarrow \mathcal{B}$ which maps every $n \in \mathbb{N} = \text{obj}(\mathcal{B}_{\otimes})$ to the single object in \mathcal{B} and sends $\alpha \in \text{hom}(n, m)$ to

$$R(\alpha) = \psi_m^{-1} \circ \alpha \circ \psi_n \in \text{Aut}(A \otimes \mathbb{K})$$

This is indeed functorial, since for $\beta \in \text{hom}(m, l)$ we have

$$R(\beta \circ \alpha) = \psi_l^{-1} \circ (\beta \circ \alpha) \circ \psi_n = (\psi_l^{-1} \circ \beta \circ \psi_m) \circ (\psi_m^{-1} \circ \alpha \circ \psi_n) = R(\beta) \circ R(\alpha) .$$

By the choice of ψ_1 the functor $R \circ J$ is the identity on \mathcal{B} . The functor $J \circ R$ maps every $n \in \mathbb{N}$ to $1 \in \mathbb{N}$ and sends $\alpha \in \text{hom}(n, m)$ to $\psi_m^{-1} \circ \alpha \circ \psi_n \in \text{hom}(1, 1)$. Therefore $\{\psi_n \in \text{hom}(1, n) : n \geq 1\}$ provides a natural transformation between $J \circ R$ and the identity functor on \mathcal{B}_\otimes .

Let $F: \mathcal{B} \times \mathcal{B} \rightarrow \mathcal{B}_\otimes$ be the functor $J \circ \otimes_\psi$ and set $G = \otimes \circ (J \times J)$. Note that $F(pt, pt) = 1$ and $G(pt, pt) = 2$, where pt denotes the single object of \mathcal{B} . Set $\kappa_{pt} = \psi \in \text{hom}(1, 2)$. It is easy to check that this choice provides a natural transformation $\kappa: F \rightarrow G$. \square

Corollary 3.7. *The space $B\text{Aut}(A \otimes \mathbb{K})_\otimes$ has a homotopy unit with respect to the tensor operation. In particular, it is a unital H -space.*

Proof. Since the diagram (2) from Lemma 3.5 is commutative up to a natural transformation, the corresponding diagram of the geometric realizations

$$\begin{array}{ccc} B\text{Aut}(A \otimes \mathbb{K}) \times B\text{Aut}(A \otimes \mathbb{K}) & \xrightarrow{\mu_\psi} & B\text{Aut}(A \otimes \mathbb{K}) \\ \downarrow & & \downarrow \\ B\text{Aut}(A \otimes \mathbb{K})_\otimes \times B\text{Aut}(A \otimes \mathbb{K})_\otimes & \longrightarrow & B\text{Aut}(A \otimes \mathbb{K})_\otimes \end{array}$$

commutes up to homotopy. From this, it is easy to deduce that the image of the basepoint in $B\text{Aut}(A \otimes \mathbb{K})_\otimes$ is a homotopy unit for the multiplication induced by \otimes . \square

So far, we have proven that $B\text{Aut}(A \otimes \mathbb{K})$ and $B\text{Aut}(A \otimes \mathbb{K})_\otimes$ are both homotopy unital H -spaces, which are both grouplike, because π_0 is trivial in both cases. Furthermore, we know that $\Omega B(|N_\bullet \mathcal{B}_\otimes^+|) = \Omega B B\text{Aut}(A \otimes \mathbb{K})_\otimes^+$ is an infinite loop space by Theorem 3.3.

Theorem 3.8. *For any strongly self-absorbing C^* -algebra A the space $B\text{Aut}(A \otimes \mathbb{K})_\otimes$ is an infinite loop space with respect to its natural tensor product operation. In particular, the monoid of isomorphism classes $(\text{Bun}_X(A \otimes \mathbb{K}), \otimes)$ of principal $\text{Aut}(A \otimes \mathbb{K})$ -bundles is an abelian group with respect to the tensor product. Furthermore, $B\text{Aut}(A \otimes \mathbb{K})_\otimes$ is the first space in a spectrum defining a cohomology theory E_A^\bullet with $E_A^0(X) = [X, \text{Aut}(A \otimes \mathbb{K})]$ and $E_A^1(X) = \text{Bun}_X(A \otimes \mathbb{K})$. If X is a compact metrizable space and $A \neq \mathbb{C}$, then $E_A^0(X) \cong K_0(C(X) \otimes A)_+^\times$.*

Proof. Applying Lemma 3.4 to the inclusion functor $\mathcal{B}_\otimes \rightarrow \mathcal{B}_\otimes^+$ we obtain that $B\text{Aut}(A \otimes \mathbb{K})_\otimes = |N_\bullet \mathcal{B}_\otimes|$ is an infinite loop space. This implies the second statement, since the H -spaces $B\text{Aut}(A \otimes \mathbb{K})$ and $B\text{Aut}(A \otimes \mathbb{K})_\otimes$ are homotopy equivalent by Lemma 3.6. As described above, an infinite loop space yields a spectrum and therefore a cohomology theory via iterated delooping. If we consider $B\text{Aut}(A \otimes \mathbb{K})_\otimes$ as the first space of the spectrum, we obtain the 0th one by forming the loop space. But this is

$$\Omega B\text{Aut}(A \otimes \mathbb{K})_\otimes \simeq \Omega B\text{Aut}(A \otimes \mathbb{K}) \simeq \text{Aut}(A \otimes \mathbb{K}),$$

which proves the third statement. The fourth statement follows from Theorem 2.22. \square

Corollary 3.9. *For any strongly self-absorbing C^* -algebra A the space $B\text{Aut}_0(A \otimes \mathbb{K})_\otimes$ is an infinite loop space with respect to its natural tensor product operation. The corresponding cohomology theory is denoted by $\bar{E}_A^*(X)$.*

Proof. The proof is entirely similar to the proof of Theorem 3.8, except that we replace the category \mathcal{B} by the topological category \mathcal{B}^0 which has as its object space just a single point and the group $\text{Aut}_0(A \otimes \mathbb{K})$ as its morphism space and we replace the category \mathcal{B}_\otimes by the category \mathcal{B}_\otimes^0 defined as follows. The object space of \mathcal{B}_\otimes^0 is \mathbb{N} . The morphisms $\text{hom}(n, m)$ are given by those maps α in $\text{Iso}((A \otimes \mathbb{K})^{\otimes n}, (A \otimes \mathbb{K})^{\otimes m})$ with the property that $[\alpha((1 \otimes e)^{\otimes n})] = [(1 \otimes e)^{\otimes m}]$ in $K_0((A \otimes \mathbb{K})^{\otimes m})$. This condition will insure that the functor $R : \mathcal{B}_\otimes^0 \rightarrow \mathcal{B}^0$ from the proof of Lemma 3.6 is well-defined. \square

Remark 3.10. We have seen that the classifying space $B\text{Aut}(A \otimes \mathbb{K})$ has the homotopy type of a CW complex. Since its homotopy groups are countable, it follows that this space is homotopy equivalent to a locally finite simplicial complex and hence to an absolute neighborhood extensor, see [28, Thm.6.1, p.137]. It follows that $E_A^1(X)$ is a continuous functor in the sense that if X is the projective limit of projective system $(X_n)_n$ of compact metrizable spaces, then $E_A^1(X) \cong \varprojlim E_A^1(X_n)$, see [20, Thm.11.9, p.287]. Since any compact metrizable space X is the projective limit of a system of finite polyhedra $(X_n)_n$ by [20], one can approach the computation of $E_A^1(X)$ by first computing $E_A^1(X_n)$ using the Atiyah-Hirzebruch spectral sequence.

4. A GENERALIZED DIXMIER-DOUADY THEORY

Definition 4.1. Let B be a continuous field C^* -algebra over a locally compact metrizable space X whose fibers are stably isomorphic to strongly self-absorbing C^* -algebras, which are not necessarily mutual isomorphic. We say that B satisfies *the Fell condition* if for each point $x \in X$, there is a closed neighborhood V of x and a projection $p \in B(V)$ such that $[p(v)] \in K_0(B(v))^\times$ for all $v \in V$. If one can choose $V = X$, then we say that B satisfies the global Fell condition.

Theorem 4.2. *Let A be a strongly self-absorbing C^* -algebra. Let X be a locally compact space of finite covering dimension and let B be a separable continuous field C^* -algebra over X with all fibers abstractly isomorphic to $A \otimes \mathbb{K}$. Then B is locally trivial if and only if it satisfies Fell's condition. If X is compact, then B is trivial if and only if B satisfies the global Fell condition.*

Proof. Suppose that there is a projection $p \in B(V)$ such that $[p(v)] \in K_0(B(v))^\times$ for all v in a compact subset V of X . We will show that $B(V) \cong C(V) \otimes A \otimes \mathbb{K}$. First we observe that by Lemma 2.14 it follows that $p(v)B(v)p(v) \cong A$, since $B(v) \cong A \otimes \mathbb{K}$. Therefore $pB(V)p$ is a unital continuous field over a finite dimensional space with fibers isomorphic to A and hence $pB(V)p \cong C(V) \otimes A$ by [15]. Second, since p is a full projection, we have that $pB(V)p \otimes \mathbb{K} \cong B(V) \otimes \mathbb{K}$ as $C(V)$ -algebras by [9]. Third, $B(V) \otimes \mathbb{K} \cong B(V)$ by [23] since V is finite dimensional and each fiber of B is stable. Putting these facts together we obtain the desired conclusion:

$$B(V) \cong B(V) \otimes \mathbb{K} \cong pB(V)p \otimes \mathbb{K} \cong C(V) \otimes A \otimes \mathbb{K}. \quad \square$$

Having obtained an efficient criterion for local triviality, we now turn to the question of classifying locally trivial continuous field C^* -algebras by cohomological invariants. Let X be a finite connected CW complex. Let $R = K_0(A)$ and let R_+^\times denote the multiplicative abelian group

$K_0(A)_+^\times$. If A is purely infinite, then $K_0(A)_+^\times = K_0(A)^\times$ and so $R_+^\times = R^\times$. Suppose that A satisfies the UCT. Then $K_1(A) = 0$ by [49].

The coefficients of the generalized cohomology theory $E_A^*(X)$ were computed in Theorem 2.18. Consequently, by [22], the E_2 -page of the Atiyah-Hirzebruch spectral sequence for the generalized cohomology $E_A^*(X)$, $A \neq \mathbb{C}$, looks as shown below.

	0	1	2	3
0	$H^0(X, R_+^\times)$	$H^1(X, R_+^\times)$	$H^2(X, R_+^\times)$	$H^3(X, R_+^\times)$
-1	0	0	0	0
-2	$H^0(X, R)$	$H^1(X, R)$	$H^2(X, R)$	$H^3(X, R)$
-3	0	0	0	0
-4	$H^0(X, R)$	$H^1(X, R)$	$H^2(X, R)$	$H^3(X, R)$

If $A = \mathbb{C}$, all the rows of the E_2 -page of $E_{\mathbb{C}}^1(X)$ are null with the exception of the (-2) -row whose entries are $H^p(X, \mathbb{Z})$, $p \geq 0$. Since the differentials in the Atiyah-Hirzebruch spectral sequence are torsion operators, [2, Thm.2.7], we obtain the following.

Corollary 4.3. *Let X be a finite connected CW complex such that $H^*(X, R)$ is torsion free. If $A \neq \mathbb{C}$ satisfies the UCT, then*

$$\mathcal{B}un_X(A \otimes \mathbb{K}) \cong E_A^1(X) \cong H^1(X, R_+^\times) \times \prod_{k \geq 1} H^{2k+1}(X, R) .$$

Corollary 4.4. *Let X be a compact metrizable space and let $M_{\mathbb{Q}}$ denote the universal UHF-algebra with $K_0(M_{\mathbb{Q}}) \cong \mathbb{Q}$. Then there are natural isomorphism of groups*

$$\mathcal{B}un_X(M_{\mathbb{Q}} \otimes \mathbb{K}) \cong E_{M_{\mathbb{Q}}}^1(X) \cong H^1(X, \mathbb{Q}_+^\times) \oplus \bigoplus_{k \geq 1} H^{2k+1}(X, \mathbb{Q}) .$$

$$\mathcal{B}un_X(M_{\mathbb{Q}} \otimes \mathcal{O}_\infty \otimes \mathbb{K}) \cong E_{M_{\mathbb{Q}} \otimes \mathcal{O}_\infty}^1(X) \cong H^1(X, \mathbb{Q}^\times) \oplus \bigoplus_{k \geq 1} H^{2k+1}(X, \mathbb{Q}) .$$

Proof. Set $h^*(X) = E_A^*(X)$, $\bar{h}^*(X) = \bar{E}_A^*(X)$ (see Cor. 3.9) and $R = K_0(A)$ where A is either $M_{\mathbb{Q}}$ or $M_{\mathbb{Q}} \otimes \mathcal{O}_\infty$. We will show that there are natural isomorphisms (i) $h^1(X) \cong H^1(X, R_+^\times) \oplus \bar{h}^1(X)$ and (ii) $\bar{h}^1(X) \cong \bigoplus_{k \geq 1} H^{2k+1}(X, \mathbb{Q})$. Note that $\bar{h}^*(pt) = t\mathbb{Q}[t]$ with $\deg(t) = -2$ and $h^*(pt) = R_+^\times \oplus t\mathbb{Q}[t]$. Suppose first that X is a finite connected CW-complex. Then (ii) follows by applying the isomorphism established in equation (3.20) of [22, p.48] since $\bar{h}^*(pt)$ is a vector spaces over \mathbb{Q} . If G is a topological group and H a normal subgroup of G such that $H \rightarrow G \rightarrow G/H$ is a principal H -bundle, then there is a homotopy fibre sequence $G \rightarrow G/H \rightarrow BH \rightarrow BG \rightarrow B(G/H)$ and hence an exact sequence of pointed sets $[X, G] \rightarrow [X, G/H] \rightarrow [X, BH] \rightarrow [X, BG] \rightarrow [X, B(G/H)]$.

Using this for the principal bundle from Corollary 2.19 we obtain an exact sequence of groups $0 \rightarrow \bar{h}^1(X) \rightarrow h^1(X) \xrightarrow{\delta_0} H^1(X, R_+^\times)$. We want to compare this sequence with the exact sequence $0 \rightarrow F^2 h^1(X) \rightarrow h^1(X) \rightarrow H^1(X, R_+^\times) \rightarrow 0$ given by the Atiyah-Hirzebruch spectral sequence. Recall that $F^2 h^1(X) = \ker(h^1(X) \rightarrow h^1(X_1))$, where X_1 is the 1-skeleton of X . Since both maps with target $H^1(X_1, R_+^\times)$ are injective in the following commutative diagram induced by $X_1 \hookrightarrow X$

$$\begin{array}{ccc} h^1(X) & \xrightarrow{\delta_0} & H^1(X, R_+^\times) \\ \downarrow & & \downarrow \\ h^1(X_1) & \longrightarrow & H^1(X_1, R_+^\times) \end{array}$$

we deduce that $F^2 h^1(X) \cong \ker(\delta_0) \cong \bar{h}^1(X)$ and hence obtain an exact sequence $0 \rightarrow \bar{h}^1(X) \rightarrow h^1(X) \rightarrow H^1(X, R_+^\times) \rightarrow 0$. Since $\bar{h}^1(X)$ is a divisible group it follows that $h^1(X)$ splits as $H^1(X, R_+^\times) \oplus \bar{h}^1(X)$. To verify that there is a natural splitting one employs the natural transformation $h^*(X) \rightarrow \bar{h}^*(X)$ induced by the coefficient map $h^*(pt) \rightarrow \bar{h}^*(pt)$, $(r, f(t)) \mapsto f(t)$, see [22, Thm.3.22].

For the general case we write X as a projective limit of a system of polyhedra $(X_n)_n$ and then we apply the continuity property of $E_A^1(X)$ as discussed in Remark 3.10. \square

Let A be a strongly self-absorbing C^* -algebra that satisfies the UCT. Then $A \otimes M_{\mathbb{Q}} \otimes \mathcal{O}_{\infty} \cong M_{\mathbb{Q}} \otimes \mathcal{O}_{\infty}$ by [49]. The canonical unital embedding $A \rightarrow A \otimes M_{\mathbb{Q}} \otimes \mathcal{O}_{\infty} \cong M_{\mathbb{Q}} \otimes \mathcal{O}_{\infty}$ induces a morphism of groups $\text{Aut}(A \otimes \mathbb{K}) \rightarrow \text{Aut}(M_{\mathbb{Q}} \otimes \mathcal{O}_{\infty} \otimes \mathbb{K})$ and hence a morphism of groups

$$\delta : E_A^1(X) \rightarrow E_{M_{\mathbb{Q}} \otimes \mathcal{O}_{\infty}}^1(X) \cong H^1(X, \mathbb{Q}^\times) \oplus \bigoplus_{k \geq 1} H^{2k+1}(X, \mathbb{Q}).$$

Definition 4.5. We define rational characteristic classes $\delta_0 : \mathcal{Bun}_X(A \otimes \mathbb{K}) \rightarrow H^1(X, \mathbb{Q}^\times)$ and $\delta_k : \mathcal{Bun}_X(A \otimes \mathbb{K}) \rightarrow H^{2k+1}(X, \mathbb{Q})$, $k \geq 1$, to be the components of the map δ from above. It is clear that δ_0 lifts to a map $\delta_0 : \mathcal{Bun}_X(A \otimes \mathbb{K}) \rightarrow H^1(X, K_0(A)_+^\times)$ induced by the morphism of groups $\text{Aut}(A \otimes \mathbb{K}) \rightarrow \pi_0(\text{Aut}(A \otimes \mathbb{K})) \cong K_0(A)_+^\times$ and which gives the obstruction to reducing the structure group to $\text{Aut}_0(A \otimes \mathbb{K})$. We will see in Corollary 4.7 that δ_1 also lifts to an integral class with values in $H^3(X, \mathbb{Z})$. One has $\delta_k(B_1 \otimes B_2) = \delta_k(B_1) + \delta_k(B_2)$, $k \geq 0$.

Since the differentials in the Atiyah-Hirzebruch spectral sequence are torsion operators we deduce:

Corollary 4.6. *Let A be a strongly self-absorbing C^* -algebra that satisfies the UCT. Let X be a finite connected CW complex such that $H^*(X, \mathbb{Z})$ is torsion free. Then:*

- (i) $B_1, B_2 \in \mathcal{Bun}_X(A \otimes \mathbb{K})$ are isomorphic if and only if $\delta_k(B_1) = \delta_k(B_2)$ for all $k \geq 0$.
- (ii) $\mathcal{Bun}_X(\mathcal{Z} \otimes \mathbb{K}) \cong \bigoplus_{k \geq 1} H^{2k+1}(X, \mathbb{Z})$ and $\mathcal{Bun}_X(\mathcal{O}_{\infty} \otimes \mathbb{K}) \cong H^1(X, \mathbb{Z}/2) \oplus \bigoplus_{k \geq 1} H^{2k+1}(X, \mathbb{Z})$.

Corollary 4.7. *Let X be a compact connected metrizable space. Then $E_{\mathbb{C}}^i(X) \cong H^{2+i}(X, \mathbb{Z})$ is a natural direct summand of both $E_{\mathcal{Z}}^i(X)$ and $E_{\mathcal{O}_{\infty}}^i(X)$ for all $i \geq 0$.*

Proof. Using the continuity properties discussed in Remark 3.10, we may assume that X is a finite connected CW complex. The unital map $\mathbb{C} \rightarrow \mathcal{Z}$ induce a natural transformation of cohomology

theories $T : E_{\mathbb{C}}^*(X) \rightarrow E_{\mathbb{Z}}^*(X)$. The desired conclusion for $E_{\mathbb{Z}}^*(X)$ follows now from the naturality of the Atiyah-Hirzebruch spectral sequence since all the rows of the E_2 -page of $E_{\mathbb{C}}^*(X)$ are null with the exception of the (-2) -row and T induces the identity map on this row due to the isomorphism $\pi_2(\text{Aut}(\mathbb{K})) \rightarrow \pi_2(\text{Aut}(\mathbb{Z} \otimes \mathbb{K}))$, see Theorem 2.18. The corresponding result for \mathcal{O}_{∞} is deduced from the result for \mathbb{Z} by using Remark 2.20 and Corollary 2.21 applied to $A = \mathbb{Z}$. \square

Corollary 4.8. *Let X be a compact metrizable space and let A be a strongly self-absorbing C^* -algebra. Two bundles $B_1, B_2 \in \mathcal{Bun}_X(A \otimes \mathbb{K})$ are isomorphic if and only if $B_1 \otimes \mathcal{O}_{\infty} \cong B_2 \otimes \mathcal{O}_{\infty}$.*

Proof. Without any loss of generality we may assume that X is a finite CW-complex. By Corollary 2.19 there is a commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Aut}_0(A \otimes \mathbb{K}) & \longrightarrow & \text{Aut}(A \otimes \mathbb{K}) & \longrightarrow & K_0(A)_+^{\times} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \text{Aut}_0(\mathcal{O}_{\infty} \otimes A \otimes \mathbb{K}) & \longrightarrow & \text{Aut}(\mathcal{O}_{\infty} \otimes A \otimes \mathbb{K}) & \longrightarrow & K_0(A)^{\times} \longrightarrow 0 \end{array}$$

Passing to classifying spaces we obtain a commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & [X, B\text{Aut}_0(A \otimes \mathbb{K})] & \longrightarrow & [X, B\text{Aut}(A \otimes \mathbb{K})] & \longrightarrow & H^1(X, K_0(A)_+^{\times}) \\ & & \downarrow i & & \downarrow T & & \downarrow j \\ 0 & \longrightarrow & [X, B\text{Aut}_0(\mathcal{O}_{\infty} \otimes A \otimes \mathbb{K})] & \longrightarrow & [X, B\text{Aut}(\mathcal{O}_{\infty} \otimes A \otimes \mathbb{K})] & \longrightarrow & H^1(X, K_0(A)^{\times}) \end{array}$$

This is a diagram of abelian groups by Theorem 3.8 and Corollary 3.9. The map j is injective. This follows from the exactness of the sequence $H^0(X, R_2) \rightarrow H^0(X, R_2/R_1) \rightarrow H^1(X, R_1) \rightarrow H^1(X, R_2)$ induced by an inclusion of discrete abelian groups $R_1 \hookrightarrow R_2$. Let us argue that the map i is also injective. If $A \neq \mathbb{C}$, this follows from Corollary 2.21, whereas for $A = \mathbb{C}$ this was proved in Corollary 4.7. The five lemma implies now that the map $T : E_A^1(X) \rightarrow E_{A \otimes \mathcal{O}_{\infty}}^1(X)$ is injective. \square

Finally we address the question to what extent the results (i) and (ii) of Dixmier and Douady mentioned in the introduction admit generalizations to our context. The following statement corresponds to (i) and the first part of (ii).

Corollary 4.9. *Let B be a separable continuous field C^* -algebra over a compact metrizable space whose fibers are Morita equivalent to the same strongly self-absorbing C^* -algebra A . Suppose that B satisfies Fell's condition. Then for each $x \in X$, there is a closed neighborhood V of x with the following property. There exists a unital separable continuous field D over V with fibers isomorphic to A such that $B(V) \otimes \mathbb{K} \cong D \otimes \mathbb{K}$. If A is finite dimensional or if X is finite dimensional, then $B \otimes \mathbb{K}$ is locally trivial and therefore we can associate to it an invariant $\delta(B) \in E_A^1(X)$ which classifies $B \otimes \mathbb{K}$ up to isomorphism of continuous fields, and B up to Morita equivalence over X .*

Proof. Let $x \in X$. Let p and V be as in Definition 4.1. Letting $D := pB(V)p$ we have already seen in the proof of Theorem 4.2 that $B(V) \otimes \mathbb{K} \cong D \otimes \mathbb{K}$ and that all the fibers of D are isomorphic to A . If A is finite dimensional, then $A = \mathbb{C}$, and so obviously $D = C(V)$. This corresponds to the

result (i) of Dixmier and Douady. If X is finite dimensional, then $B(V) \otimes \mathbb{K} \cong C(V) \otimes A \otimes \mathbb{K}$ by Theorem 4.2. We conclude the proof by applying Theorem 3.8. \square

As we have just seen, the class $\delta(B) \in E_A^1(X)$ is defined for continuous fields with fibers Morita equivalent to A which satisfy Fell's condition. Furthermore, one can associate rational characteristic classes to certain continuous fields which are typically very far from being locally trivial and whose fibers are not necessarily Morita equivalent to each other.

Corollary 4.10. *Let B be a separable continuous field C^* -algebra over a finite dimensional compact metrizable space whose fibers are Morita equivalent to (possibly different) strongly self-absorbing C^* -algebras satisfying the UCT. Suppose that B satisfies Fell's condition. Then $B \otimes \mathcal{O}_\infty \otimes M_{\mathbb{Q}} \otimes \mathbb{K}$ is locally trivial and so we can associate “stable” rational characteristic classes to B , by defining $\delta_k^{\text{stable}}(B) = \delta_k(B \otimes \mathcal{O}_\infty \otimes M_{\mathbb{Q}} \otimes \mathbb{K}) \in H^{2k+1}(X, \mathbb{Q})$.*

Proof. The fibers $B(x) \otimes \mathcal{O}_\infty \otimes M_{\mathbb{Q}} \otimes \mathbb{K}$ satisfy the UCT and they are stabilized strongly self-absorbing Kirchberg algebras. It follows by [49] that they are all isomorphic to $\mathcal{O}_\infty \otimes M_{\mathbb{Q}} \otimes \mathbb{K}$. Now $B \otimes \mathcal{O}_\infty \otimes M_{\mathbb{Q}} \otimes \mathbb{K}$ still satisfies Fell's condition and hence it is locally trivial by Theorem 4.2. We conclude the proof by applying Corollary 4.4. \square

REFERENCES

- [1] John Frank Adams. *Infinite loop spaces*, volume 90 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, N.J., 1978. 14
- [2] Dominique Arlettaz. The order of the differentials in the Atiyah-Hirzebruch spectral sequence. *K-Theory*, 6(4):347–361, 1992. 21
- [3] Michael Atiyah and Graeme Segal. Twisted K -theory. *Ukr. Mat. Visn.*, 1(3):287–330, 2004. 2
- [4] Michael Atiyah and Graeme Segal. Twisted K -theory and cohomology. In *Inspired by S. S. Chern*, volume 11 of *Nankai Tracts Math.*, pages 5–43. World Sci. Publ., Hackensack, NJ, 2006. 2
- [5] Jean Bellissard. K -theory of C^* -algebras in solid state physics. In *Statistical mechanics and field theory: mathematical aspects (Groningen, 1985)*, volume 257 of *Lecture Notes in Phys.*, pages 99–156. Springer, Berlin, 1986. 1
- [6] David J. Benson, Alex Kumjian, and N. Christopher Phillips. Symmetries of Kirchberg algebras. *Canad. Math. Bull.*, 46(4):509–528, 2003. 12
- [7] B. Blackadar. *Operator algebras*, volume 122 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 2006. Theory of C^* -algebras and von Neumann algebras, Operator Algebras and Non-commutative Geometry, III. 8
- [8] Peter Bouwknegt and Varghese Mathai. D-branes, B -fields and twisted K -theory. *J. High Energy Phys.*, (3):Paper 7, 11, 2000. 2
- [9] Lawrence G. Brown. Stable isomorphism of hereditary subalgebras of C^* -algebras. *Pacific J. Math.*, 71(2):335–348, 1977. 9, 20
- [10] Lawrence G. Brown, Philip Green, and Marc A. Rieffel. Stable isomorphism and strong Morita equivalence of C^* -algebras. *Pacific J. Math.*, 71(2):349–363, 1977. 11
- [11] Alain Connes. *Noncommutative geometry*. Academic Press Inc., San Diego, CA, 1994. 1

- [12] Joachim Cuntz and Nigel Higson. Kuiper's theorem for Hilbert modules. In *Operator algebras and mathematical physics (Iowa City, Iowa, 1985)*, volume 62 of *Contemp. Math.*, pages 429–435. Amer. Math. Soc., Providence, RI, 1987. 8
- [13] Marius Dadarlat. The homotopy groups of the automorphism group of Kirchberg algebras. *J. Noncommut. Geom.*, 1(1):113–139, 2007. 9, 13
- [14] Marius Dadarlat. Fiberwise KK -equivalence of continuous fields of C^* -algebras. *J. K-Theory*, 3(2):205–219, 2009. 3, 4
- [15] Marius Dadarlat and Wilhelm Winter. Trivialization of $C(X)$ -algebras with strongly self-absorbing fibres. *Bull. Soc. Math. France*, 136(4):575–606, 2008. 20
- [16] Marius Dadarlat and Wilhelm Winter. On the KK -theory of strongly self-absorbing C^* -algebras. *Math. Scand.*, 104(1):95–107, 2009. 2, 5
- [17] Jacques Dixmier and Adrien Douady. Champs continus d'espaces hilbertiens et de C^* -algèbres. *Bull. Soc. Math. France*, 91:227–284, 1963. 2, 3
- [18] P. Donovan and M. Karoubi. Graded Brauer groups and K -theory with local coefficients. *Inst. Hautes Études Sci. Publ. Math.*, (38):5–25, 1970. 2
- [19] B. Eckmann and P. J. Hilton. Group-like structures in general categories. I. Multiplications and comultiplications. *Math. Ann.*, 145:227–255, 1961/1962. 10
- [20] Samuel Eilenberg and Norman Steenrod. *Foundations of algebraic topology*. Princeton University Press, Princeton, New Jersey, 1952. 20
- [21] Richard H. Herman and Jonathan Rosenberg. Norm-close group actions on C^* -algebras. *J. Operator Theory*, 6(1):25–37, 1981. 7
- [22] Peter Hilton. *General cohomology theory and K-theory*, volume 1 of *Course given at the University of São Paulo in the summer of 1968 under the auspices of the Instituto de Pesquisas Matemáticas, Universidade de São Paulo. London Mathematical Society Lecture Note Series*. Cambridge University Press, London, 1971. 21, 22
- [23] Ilan Hirshberg, Mikael Rørdam, and Wilhelm Winter. $C_0(X)$ -algebras, stability and strongly self-absorbing C^* -algebras. *Math. Ann.*, 339(3):695–732, 2007. 3, 4, 20
- [24] Peter J. Huber. Homotopical cohomology and Čech cohomology. *Math. Ann.*, 144:73–76, 1961. 2
- [25] Xinhui Jiang. Nonstable K -theory for \mathcal{Z} -stable C^* -algebras. arXiv:9707.5228[math.OA], 1997. 5, 12
- [26] Kazunori Kodaka. Full projections, equivalence bimodules and automorphisms of stable algebras of unital C^* -algebras. *J. Operator Theory*, 37(2):357–369, 1997. 9, 11
- [27] N. P. Landsman. *Mathematical topics between classical and quantum mechanics*. Springer Monographs in Mathematics. Springer-Verlag, New York, 1998. 1
- [28] A.T. Lundell and S. Weingram. *The topology of CW complexes*. The University Series in Higher Mathematics. New York etc.: Van Nostrand Reinhold Company. VIII, 216 p. , 1969. 8, 20
- [29] J. P. May. *The geometry of iterated loop spaces*. Springer-Verlag, Berlin, 1972. Lectures Notes in Mathematics, Vol. 271. 14, 15
- [30] J. P. May. E_∞ spaces, group completions, and permutative categories. In *New developments in topology (Proc. Sympos. Algebraic Topology, Oxford, 1972)*, pages 61–93. London Math. Soc. Lecture Note Ser., No. 11. Cambridge Univ. Press, London, 1974. 15
- [31] J. P. May. The spectra associated to permutative categories. *Topology*, 17(3):225–228, 1978. 14, 15
- [32] J. Peter May. Classifying spaces and fibrations. *Mem. Amer. Math. Soc.*, 1(1, 155):xiii+98, 1975. 17

- [33] D. McDuff and G. Segal. Homology fibrations and the “group-completion” theorem. *Invent. Math.*, 31(3):279–284, 1975/76. 16
- [34] J. A. Mingo. K -theory and multipliers of stable C^* -algebras. *Trans. Amer. Math. Soc.*, 299(1):397–411, 1987. 9, 10
- [35] Norio Nawata and Yasuo Watatani. Fundamental group of simple C^* -algebras with unique trace. *Adv. Math.*, 225(1):307–318, 2010. 11
- [36] Victor Nistor. Fields of AF-algebras. *J. Operator Theory*, 28(1):3–25, 1992. 2
- [37] Marc A. Rieffel. Quantization and C^* -algebras. In *C^* -algebras: 1943–1993 (San Antonio, TX, 1993)*, volume 167 of *Contemp. Math.*, pages 66–97. Amer. Math. Soc., Providence, RI, 1994. 1
- [38] M. Rørdam. *Classification of nuclear, simple C^* -algebras*, volume 126 of *Encyclopaedia Math. Sci.* Springer, Berlin, 2002. 11, 13
- [39] Mikael Rørdam. The stable and the real rank of \mathcal{Z} -absorbing C^* -algebras. *Internat. J. Math.*, 15(10):1065–1084, 2004. 5, 13
- [40] Jonathan Rosenberg. Continuous-trace algebras from the bundle theoretic point of view. *J. Austral. Math. Soc. Ser. A*, 47(3):368–381, 1989. 2
- [41] Claude Schochet. The Dixmier-Douady invariant for dummies. *Notices Amer. Math. Soc.*, 56(7):809–816, 2009. 2
- [42] Rolf Schön. Fibrations over a CWh-base. *Proc. Amer. Math. Soc.*, 62(1):165–166 (1977), 1976. 8
- [43] Graeme Segal. Classifying spaces and spectral sequences. *Inst. Hautes Études Sci. Publ. Math.*, (34):105–112, 1968. 18
- [44] Graeme Segal. Categories and cohomology theories. *Topology*, 13:293–312, 1974. 4, 14, 15, 16
- [45] Graeme Segal. Topological structures in string theory. *R. Soc. Lond. Philos. Trans. Ser. A Math. Phys. Eng. Sci.*, 359(1784):1389–1398, 2001. Topological methods in the physical sciences (London, 2000). 2
- [46] Arne Strøm. Note on cofibrations. *Math. Scand.*, 19:11–14, 1966. 14
- [47] Robert W. Thomason. First quadrant spectral sequences in algebraic K -theory via homotopy colimits. *Comm. Algebra*, 10(15):1589–1668, 1982. 15
- [48] Klaus Thomsen. Homotopy classes of $*$ -homomorphisms between stable C^* -algebras and their multiplier algebras. *Duke Math. J.*, 61(1):67–104, 1990. 9
- [49] Andrew S. Toms and Wilhelm Winter. Strongly self-absorbing C^* -algebras. *Trans. Amer. Math. Soc.*, 359(8):3999–4029, 2007. 2, 4, 5, 9, 12, 21, 22, 24
- [50] Wilhelm Winter. Strongly self-absorbing C^* -algebras are \mathcal{Z} -stable. *J. Noncommut. Geom.*, 5(2):253–264, 2011. 2, 5

MD: DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, WEST LAFAYETTE, IN 47907, USA
E-mail address: mdd@math.purdue.edu

UP: MATHEMATISCHES INSTITUT, WESTFÄLISCHE WILHELMS-UNIVERSITÄT MÜNSTER, EINSTEINSTRASSE 62,
 48149 MÜNSTER, GERMANY
E-mail address: u.pennig@uni-muenster.de